

Sptrx-2 (N-14): sc-160842

BACKGROUND

Sptrx-2 (spermatid-specific thioredoxin-2), also known as NME8, CILD6, SPTRX2 or TXNDC3 (thioredoxin domain-containing protein 3), is a 588 amino acid cytoplasmic and testis-specific protein belonging to the NDK family. Expressed only in primary spermatocytes and round spermatids, Sptrx-2 may be required during the final stages of sperm tail maturation in the testis and/or epididymis, where extensive disulfide bonding of fibrous sheath (FS) proteins occur. Sptrx-2 contains a thioredoxin domain and three inactive NDK domains that each lack the active His residue, suggesting that they are not capable of NDP kinase activity. Defects in the gene encoding Sptrx-2 are the cause of primary ciliary dyskinesia type 6, an autosomal recessive disorder characterized by axonemal abnormalities of motile cilia. Primary ciliary dyskinesia associated with situs inversus is referred to as Kartagener syndrome.

REFERENCES

1. Sadek, C.M., et al. 2001. Sptrx-2, a fusion protein composed of one thioredoxin and three tandemly repeated NDP-kinase domains is expressed in human testis germ cells. *Genes Cells* 6: 1077-1090.
2. Mahr, S., et al. 2006. Cis- and transacting gene regulation is associated with osteoarthritis. *Am. J. Hum. Genet.* 78: 793-803.
3. Loughlin, J., et al. 2007. Genetic association analysis of RHOB and TXNDC3 in osteoarthritis. *Am. J. Hum. Genet.* 80: 383-386.
4. Duriez, B., et al. 2007. A common variant in combination with a nonsense mutation in a member of the thioredoxin family causes primary ciliary dyskinesia. *Proc. Natl. Acad. Sci. USA* 104: 3336-3341.
5. Shi, D., et al. 2008. Association of single-nucleotide polymorphisms in RHOB and TXNDC3 with knee osteoarthritis susceptibility: two case-control studies in East Asian populations and a meta-analysis. *Arthritis Res. Ther.* 10: R54.
6. Geremek, M., et al. 2008. Sequence analysis of 21 genes located in the Kartagener syndrome linkage region on chromosome 15q. *Eur. J. Hum. Genet.* 16: 688-695.
7. Faily, M., et al. 2008. DNAI1 mutations explain only 2% of primary ciliary dyskinesia. *Respiration* 76: 198-204.
8. Faily, M., et al. 2009. Mutations in DNAH5 account for only 15% of a non-preselected cohort of patients with primary ciliary dyskinesia. *J. Med. Genet.* 46: 281-286.
9. Escudier, E., et al. 2009. Ciliary defects and genetics of primary ciliary dyskinesia. *Paediatr. Respir. Rev.* 10: 51-54.

CHROMOSOMAL LOCATION

Genetic locus: TXNDC3 (human) mapping to 7p14.1; Txndc3 (mouse) mapping to 13 A2.

SOURCE

Sptrx-2 (N-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of Sptrx-2 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-160842 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Sptrx-2 (N-14) is recommended for detection of Sptrx-2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with Sptrx-1 or Sptrx-3.


Sptrx-2 (N-14) is also recommended for detection of Sptrx-2 in additional species, including canine.

Suitable for use as control antibody for Sptrx-2 siRNA (h): sc-89740, Sptrx-2 siRNA (m): sc-153807, Sptrx-2 shRNA Plasmid (h): sc-89740-SH, Sptrx-2 shRNA Plasmid (m): sc-153807-SH, Sptrx-2 shRNA (h) Lentiviral Particles: sc-89740-V and Sptrx-2 shRNA (m) Lentiviral Particles: sc-153807-V.

Molecular Weight of Sptrx-2: 67 kDa.

Positive Controls: Sptrx-2 (h2): 293T Lysate : sc-158977.

DATA



Sptrx-2 (N-14): sc-160842. Western blot analysis of Sptrx-2 expression in non-transfected: sc-117752 (A) and human Sptrx-2 transfected: sc-158977 (B) 293T whole cell lysates.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **Sptrx-2 (KK-M5): sc-135567**, our highly recommended monoclonal alternative to Sptrx-2 (N-14).