

# TTC1 (P-20): sc-160882

## BACKGROUND

The tetratricopeptide repeat (TPR) motif is a degenerate, 34 amino acid sequence found in many proteins and acts to mediate protein-protein interactions in various pathways. At the sequence level, there can be up to 16 tandem TPR repeats, each of which has a helix-turn-helix shape that stacks on other TPR repeats to achieve ligand binding specificity. TTC1 (tetratricopeptide repeat domain 1), also known as TPR1, is a 292 amino acid protein containing three TPR repeats. Considered a chaperone adaptor, TTC1 regulates HSP 70-dependent folding processes by interacting with the C-terminal domain of HSP 70. TTC1 also interacts with Ras and a few G $\alpha$  proteins, suggesting a function in protein-protein interaction relating to G-protein signaling. The gene encoding TTC1 is located on human chromosome 5, which contains 181 million base pairs and comprises nearly 6% of the human genome.

## REFERENCES

- Young, J.C., Obermann, W.M. and Hartl, F.U. 1998. Specific binding of tetratricopeptide repeat proteins to the C-terminal 12-kDa domain of hsp90. *J. Biol. Chem.* 273: 18007-18010.
- Marty, C., Browning, D.D. and Ye, R.D. 2003. Identification of tetratricopeptide repeat 1 as an adaptor protein that interacts with heterotrimeric G proteins and the small GTPase Ras. *Mol. Cell. Biol.* 23: 3847-3858.
- Oh, W.K. and Song, J. 2003. Cooperative interaction of Hsp40 and TPR1 with Hsp70 reverses Hsp70-HspBp1 complex formation. *Mol. Cells.* 16: 84-91.
- Cortajarena, A.L., Kajander, T., Pan, W., Cocco, M.J. and Regan, L. 2004. Protein design to understand peptide ligand recognition by tetratricopeptide repeat proteins. *Protein Eng. Des. Sel.* 17: 399-409.
- Kaneko, A., Umeyama, T., Utena-Abe, Y., Yamagoe, S., Niimi, M. and Uehara, Y. 2006. Tcc1p, a novel protein containing the tetratricopeptide repeat motif, interacts with Tup1p to regulate morphological transition and virulence in *Candida albicans*. *Eukaryotic Cell.* 5: 1894-1905.
- Morohashi, H., Maculins, T. and Labib, K. 2009. The amino-terminal TPR domain of Dia2 tethers SCF(Dia2) to the replisome progression complex. *Curr. Biol.* 19: 1943-1949.
- Suizu, F., Hiramaki, Y., Okumura, F., Matsuda, M., Okumura, A.J., Hirata, N., Narita, M., Kohno, T., Yokota, J., Bohgaki, M., Obuse, C., Hatakeyama, S., Obata, T. and Noguchi, M. 2009. The E3 ligase TTC3 facilitates ubiquitination and degradation of phosphorylated Akt. *Dev. Cell.* 17: 800-810.
- Lin, Z., Ho, C.W. and Grierson, D. 2009. AtTRP1 encodes a novel TPR protein that interacts with the ethylene receptor ERS1 and modulates development in *Arabidopsis*. *J. Exp. Bot.* 60: 3697-3714.
- Zhang, Z., Roe, S.M., Diogon, M., Kong, E., El Alaoui, H. and Barford, D. 2010. Molecular structure of the N-terminal domain of the APC/C subunit Cdc27 reveals a homo-dimeric tetratricopeptide repeat architecture. *J. Mol. Biol.* E-published.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## CHROMOSOMAL LOCATION

Genetic locus: Ttc1 (mouse) mapping to 11 B1.1.

## SOURCE

TTC1 (P-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of TTC1 of mouse origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-160882 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

TTC1 (P-20) is recommended for detection of TTC1 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other TTC family members.

Suitable for use as control antibody for TTC1 siRNA (m): sc-154749, TTC1 shRNA Plasmid (m): sc-154749-SH and TTC1 shRNA (m) Lentiviral Particles: sc-154749-V.

Molecular Weight of TTC1: 34 kDa.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.