

## ZnT-5 (S-13): sc-161274

### BACKGROUND

Zinc, an essential element required for cell proliferation and differentiation, plays a role in a diverse array of cellular functions (such as neuroregulation) and acts as a cofactor for numerous enzymes and transcription factors. The zinc transporter (ZnT) family regulates the supply of zinc within cells, and its members commonly contain six membrane-spanning domains, a large histidine-rich intracellular loop and a C-terminal tail. ZnT-5 (zinc transporter 5), also known as SLC30A5 (solute carrier family 30 member 5), ZNTL1 or ZTL1, is a 765 amino acid protein that localizes to the membrane of the *trans*-Golgi network. Expressed throughout the body with highest expression in liver, pancreas and kidney, ZnT-5 functions as zinc transporter that regulates zinc homeostasis within vesicular compartments and the Golgi apparatus and may help to form Insulin crystals within pancreatic  $\beta$  cells. ZnT-5 is expressed as two isoforms due to alternative splicing events and its expression is upregulated in response to zinc depletion.

### REFERENCES

1. Kambe, T., Narita, H., Yamaguchi-Iwai, Y., Hirose, J., Amano, T., Sugiura, N., Sasaki, R., Mori, K., Iwanaga, T. and Nagao, M. 2002. Cloning and characterization of a novel mammalian zinc transporter, zinc transporter 5, abundantly expressed in pancreatic  $\beta$  cells. *J. Biol. Chem.* 277: 19049-19055.
2. Cragg, R.A., Christie, G.R., Phillips, S.R., Russi, R.M., Küry, S., Mathers, J.C., Taylor, P.M. and Ford, D. 2002. A novel zinc-regulated human zinc transporter, hZTL1, is localized to the enterocyte apical membrane. *J. Biol. Chem.* 277: 22789-22797.
3. Devergnas, S., Chimienti, F., Naud, N., Pennequin, A., Coquerel, Y., Chantegrel, J., Favier, A. and Seve, M. 2004. Differential regulation of zinc efflux transporters ZnT-1, ZnT-5 and ZnT-7 gene expression by zinc levels: a real-time RT-PCR study. *Biochem. Pharmacol.* 68: 699-709.
4. Cragg, R.A., Phillips, S.R., Piper, J.M., Varma, J.S., Campbell, F.C., Mathers, J.C. and Ford, D. 2005. Homeostatic regulation of zinc transporters in the human small intestine by dietary zinc supplementation. *Gut* 54: 469-478.
5. Suzuki, T., Ishihara, K., Migaki, H., Matsuura, W., Kohda, A., Okumura, K., Nagao, M., Yamaguchi-Iwai, Y. and Kambe, T. 2005. Zinc transporters, ZnT5 and ZnT7, are required for the activation of alkaline phosphatases, zinc-requiring enzymes that are glycosylphosphatidylinositol-anchored to the cytoplasmic membrane. *J. Biol. Chem.* 280: 637-643.
6. Ishihara, K., Yamazaki, T., Ishida, Y., Suzuki, T., Oda, K., Nagao, M., Yamaguchi-Iwai, Y. and Kambe, T. 2006. Zinc transport complexes contribute to the homeostatic maintenance of secretory pathway function in vertebrate cells. *J. Biol. Chem.* 281: 17743-17750.

### CHROMOSOMAL LOCATION

Genetic locus: SLC30A5 (human) mapping to 5q13.1; Slc30a5 (mouse) mapping to 13 D1.

### SOURCE

ZnT-5 (S-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within a cytoplasmic domain of ZnT-5 of human origin.

### PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-161274 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

### APPLICATIONS

ZnT-5 (S-13) is recommended for detection of ZnT-5 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other ZnT family members.

ZnT-5 (S-13) is also recommended for detection of ZnT-5 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for ZnT-5 siRNA (h): sc-91622, ZnT-5 siRNA (m): sc-155978, ZnT-5 shRNA Plasmid (h): sc-91622-SH, ZnT-5 shRNA Plasmid (m): sc-155978-SH, ZnT-5 shRNA (h) Lentiviral Particles: sc-91622-V and ZnT-5 shRNA (m) Lentiviral Particles: sc-155978-V.

Molecular Weight (predicted) of ZnT-5 isoforms 1/2: 84/57 kDa.

Molecular Weight (observed) of ZnT-5: 68 kDa.

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

### STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.



Try **ZnT-5 (2F10-2B8): sc-517038**, our highly recommended monoclonal alternative to ZnT-5 (S-13).