

ACSVL1 (F-12): sc-161310

BACKGROUND

Acyl-coenzyme A synthetases (ACs) are a large family of related enzymes known to catalyze the fundamental initial reaction in fatty acid metabolism. The ACS family is roughly characterized based on fatty acid chain length preference among different members. The nomenclature in the ACS family reflects this relationship and includes short-chain ACS (ACSS), medium-chain ACS (ACSM), long-chain ACS (ACSL) and very long-chain ACS (ACSVL). ACSVL family members are capable of activating both long-chain fatty acids (LCFAs) and very long-chain (VLCFAs) fatty acids. There are six members of the human ACSVL subfamily which have been described as solute carrier family 27A (SLC27A) gene products. They represent a group of evolutionarily conserved fatty acid transport proteins (FATPs) recognized for their role in facilitating translocation of long-chain fatty acids across the plasma membrane. The family nomenclature has recently been unified with their respective acyl-CoA synthetase family designations: ACSVL1 (FATP2), ACSVL2 (FATP6), ACSVL3 (FATP3), ACSVL4 (FATP4), ACSVL5 (FATP1) and ACSVL6 (FATP5). ACSVLs have unique expression patterns and are found in major organs of fatty acid metabolism, such as adipose tissue, liver, heart and kidney.

REFERENCES

- Schaffer, J.E., et al. 1994. Expression cloning and characterization of a novel adipocyte long chain fatty acid transport protein. *Cell* 79: 427-436.
- Hirsch, D., et al. 1998. A family of fatty acid transporters conserved from mycobacterium to man. *Proc. Natl. Acad. Sci. USA* 95: 8625-8629.
- Abumrad, N., et al. 1999. Membrane proteins implicated in long-chain fatty acid uptake by mammalian cells: CD36, FATP, FABPm. *Biochim. Biophys. Acta* 1441: 4-13.
- Martin, G., et al. 2000. The human fatty acid transport protein-1 (SLC27A1; FATP-1) cDNA and gene: organization, chromosomal localization, and expression. *Genomics* 66: 296-304.
- Binnert, C., et al. 2000. Fatty acid transport protein-1 mRNA expression in skeletal muscle and in adipose tissue in humans. *Am. J. Physiol. Endocrinol. Metab.* 279: E1072-E1079.
- Martin, G., et al. 2000. Induction of the fatty acid transport protein 1 and acyl-CoA synthase genes by dimer-selective rexinoids suggests that the peroxisome proliferator-activated receptor-retinoid X receptor heterodimer is their molecular target. *J. Biol. Chem.* 275: 12612-12618.
- Watkins, P.A., et al. 2007. Evidence for 26 distinct acyl-coenzyme A synthetase genes in the human genome. *J. Lipid Res.* 48: 2736-2750.

CHROMOSOMAL LOCATION

Genetic locus: SLC27A2 (human) mapping to 15q21.2; Slc27a2 (mouse) mapping to 2 F1.

SOURCE

ACSVL1 (F-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping within a cytoplasmic domain of ACSVL1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-161310 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

ACSVL1 (F-12) is recommended for detection of ACSVL1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other ACSVL family members.

Suitable for use as control antibody for ACSVL1 siRNA (h): sc-90127, ACSVL1 siRNA (m): sc-140837, ACSVL1 shRNA Plasmid (h): sc-90127-SH, ACSVL1 shRNA Plasmid (m): sc-140837-SH, ACSVL1 shRNA (h) Lentiviral Particles: sc-90127-V and ACSVL1 shRNA (m) Lentiviral Particles: sc-140837-V.

Molecular Weight of ACSVL1: 70 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

SELECT PRODUCT CITATIONS

- McIntosh, A.L., et al. 2013. Liver fatty acid binding protein gene-ablation exacerbates weight gain in high-fat fed female mice. *Lipids* 48: 435-448.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **ACSVL1 (D-7): sc-393906**, our highly recommended monoclonal alternative to ACSVL1 (F-12).