

RecQL5 (N-20): sc-16157

BACKGROUND

In humans, the RecQ helicase family includes WRN, BLM, RecQL1, RecQL4 and RecQL5 proteins, all of which contain a conserved helicase domain. WRN and BLM have been demonstrated to be the responsible genes in Werner and Bloom syndromes, respectively. RecQL1 and RecQL5 also belong to the human RecQ helicase family, but their correlation with genetic disorders, if any, is unknown. The gene encoding human RecQL4, which maps to chromosome 8q24.3, is believed to be the gene responsible for the development of Rothmund-Thomson syndrome (RTS). The levels of WRN, BLM, RecQL1, RecQL4 and RecQL5 are differentially up-regulated to guarantee genomic stability in cells that are transformed or actively proliferating. In humans, RecQL1 and RecQL5 map to chromosome 12p12 and 17q25, respectively.

REFERENCES

1. Seki, M., et al. 1994. Molecular cloning of cDNA encoding human DNA helicase Q1 which has homology to *Escherichia coli* Rec Q helicase and localization of the gene at chromosome 12p12. *Nucleic Acids Res.* 22: 4566-4573.
2. Kitao, S., et al. 1999. Mutations in RecQL4 cause a subset of cases of Rothmund-Thomson syndrome. *Nat. Genet.* 22: 82-84.
3. Ohhata, T., et al. 2000. Cloning, genomic structure and chromosomal localization of the gene encoding mouse DNA helicase RecQ helicase protein-like 4. *Gene* 261: 251-258.
4. Kawabe, T., et al. 2000. Differential regulation of human RecQ family helicases in cell transformation and cell cycle. *Oncogene* 19: 4764-4772.
5. Kawabe, Y., et al. 2000. Covalent modification of the Werner's syndrome gene product with the ubiquitin-related protein, SUMO-1. *J. Biol. Chem.* 275: 20963-20966.

CHROMOSOMAL LOCATION

Genetic locus: RECQL5 (human) mapping to 17q25.2-q25.3; Recql5 (mouse) mapping to 11 E2.

SOURCE

RecQL5 (N-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of RecQL5 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-16157 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

RecQL5 (N-20) is recommended for detection of RecQL5 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for RecQL5 siRNA (h): sc-38221, RecQL5 siRNA (m): sc-38222, RecQL5 shRNA Plasmid (h): sc-38221-SH, RecQL5 shRNA Plasmid (m): sc-38222-SH, RecQL5 shRNA (h) Lentiviral Particles: sc-38221-V and RecQL5 shRNA (m) Lentiviral Particles: sc-38222-V.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **RecQL5 (C-9): sc-515050**, our highly recommended monoclonal alternative to RecQL5 (N-20).