

Josephin-3 (S-14): sc-161754

BACKGROUND

Josephin-3 (TATA box-binding protein-associated factor RNA polymerase I subunit D) is a 278 amino acid transcription factor encoded by the human TAF1D gene. The TAF1D gene product is a major component of a TBP and TAF's (TATA box-binding protein-associated factors) complex known as SL1. This SL1 complex is important for the assembly of the preinitiation complex required for RNA polymerase I-dependent transcription. The SL1/TIF-IB complex has been shown to stabilize the nucleolar transcription factor 1/UBTF on rDNA. Impaired function of the SL1 complex leads to reduced levels of Pol I transcription. The TAF1D gene product has been shown to be involved in cell cycle progression and exhibits G₂/M phase specific phosphorylations. The TAF1D gene product is expressed in most tissues and is localized to the nucleus.

REFERENCES

- Rudloff, U., Eberhard, D., Tora, L., Stunnenberg, H. and Grummt, I. 1994. TBP-associated factors interact with DNA and govern species specificity of RNA polymerase I transcription. *EMBO J.* 13: 2611-2616.
- Geiduschek, E.P. and Kassavetis, G.A. 1995. Comparing transcriptional initiation by RNA polymerases I and III. *Curr. Opin. Cell Biol.* 7: 344-351.
- Servant, N., Marcantonio, D., Th'ng, J.P. and Chalifour, L.E. 2004. TBP-associated factor 1 overexpression induces tolerance to Doxorubicin in confluent H9c2 cells by an increase in Cdk2 activity and cyclin E expression. *Mol. Cell. Biochem.* 259: 71-81.
- Friedrich, J.K., Panov, K.I., Cabart, P., Russell, J. and Zomerdijk, J.C. 2005. TBP-TAF complex SL1 directs RNA polymerase I pre-initiation complex formation and stabilizes upstream binding factor at the rDNA promoter. *J. Biol. Chem.* 280: 29551-29558.
- Gorski, J.J., Pathak, S., Panov, K., Kasciukovic, T., Panova, T., Russell, J. and Zomerdijk, J.C. 2007. A novel TBP-associated factor of SL1 functions in RNA polymerase I transcription. *EMBO J.* 26: 1560-1568.
- Kimura, J., Nguyen, S.T., Liu, H., Taira, N., Miki, Y. and Yoshida, K. 2008. A functional genome-wide RNAi screen identifies TAF1 as a regulator for apoptosis in response to genotoxic stress. *Nucleic Acids Res.* 36: 5250-5259.
- Pijnappel, W.P., Kolkman, A., Baltissen, M.P., Heck, A.J. and Timmers, H.M. 2009. Quantitative mass spectrometry of TATA binding protein-containing complexes and subunit phosphorylations during the cell cycle. *Proteome Sci.* 7: 46.
- Mayya, V., Lundgren, D.H., Hwang, S.I., Rezaul, K., Wu, L., Eng, J.K., Rodionov, V. and Han, D.K. 2009. Quantitative phosphoproteomic analysis of T cell receptor signaling reveals system-wide modulation of protein-protein interactions. *Sci. Signal.* 2: ra46.

CHROMOSOMAL LOCATION

Genetic locus: Taf1d (mouse) mapping to 9 A2.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

SOURCE

Josephin-3 (S-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Josephin-3 of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-161754 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Josephin-3 (S-14) is recommended for detection of Josephin-3 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with Josephin-1 or Josephin-2.

Suitable for use as control antibody for Josephin-3 siRNA (m): sc-146330, Josephin-3 shRNA Plasmid (m): sc-146330-SH and Josephin-3 shRNA (m) Lentiviral Particles: sc-146330-V.

Molecular Weight of Josephin-3: 32 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.