



MacroH2A1 (K-14): sc-161813

BACKGROUND

Eukaryotic histones are water soluble, basic nuclear proteins that form hetero-octameric nucleosome particles by wrapping 146 base pairs of DNA in a left-handed super-helical turn sequentially to form chromosomal fiber. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form the octamer. The octamer consists of two H2A-H2B dimers and two H3-H4 dimers, forming two nearly symmetrical halves by tertiary structure. MacroH2A1, also known as H2AFY (H2A histone family, member Y), MacroH2A1.2, MacroH2A1.1, H2A/y, H2AFJ or mH2A1, is a 372 amino acid ubiquitously expressed nuclear histone variant that is enriched in inactive X chromosome chromatin and senescence-associated heterochromatin. Involved in augmentation of signal-regulated transcription, MacroH2A1 exists as three alternatively spliced isoforms, contains one histone H2A domain and a single Macro domain.

REFERENCES

- Pehrson, J.R. and Fried, V.A. 1992. MacroH2A, a core histone containing a large nonhistone region. *Science* 257: 1398-1400.
- Pehrson, J.R., Costanzi, C. and Dharia, C. 1997. Developmental and tissue expression patterns of histone MacroH2A1 subtypes. *J. Cell. Biochem.* 65: 107-113.
- Mermoud, J.E., Costanzi, C., Pehrson, J.R. and Brockdorff, N. 1999. Histone MacroH2A1.2 relocates to the inactive X chromosome after initiation and propagation of X-inactivation. *J. Cell Biol.* 147: 1399-1408.
- Ma, Y., Jacobs, S.B., Jackson-Grusby, L., Mastrangelo, M.A., Torres-Betancourt, J.A., Jaenisch, R. and Rasmussen, T.P. 2005. DNA CpG hypomethylation induces heterochromatin reorganization involving the histone variant MacroH2A. *J. Cell. Sci.* 118: 1607-1616.
- Chakravarthy, S., Gundimella, S.K., Caron, C., Perche, P.Y., Pehrson, J.R., Khochbin, S. and Luger, K. 2005. Structural characterization of the histone variant MacroH2A. *Mol. Cell. Biol.* 25: 7616-7624.
- Kustatscher, G., Hothorn, M., Pugieux, C., Scheffzek, K. and Ladurner, A.G. 2005. Splicing regulates NAD metabolite binding to histone MacroH2A. *Nat. Struct. Mol. Biol.* 12: 624-625.
- Chu, F., Nusinow, D.A., Chalkley, R.J., Plath, K., Panning, B. and Burlingame, A.L. 2006. Mapping post-translational modifications of the histone variant MacroH2A1 using tandem mass spectrometry. *Mol. Cell Proteomics* 5: 194-203.

CHROMOSOMAL LOCATION

Genetic locus: H2AFY (human) mapping to 5q31.1; H2afy (mouse) mapping to 13 B1.

SOURCE

MacroH2A1 (K-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of MacroH2A1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-161813 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-161813 X, 200 µg/0.1 ml.

APPLICATIONS

MacroH2A1 (K-14) is recommended for detection of MacroH2A1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with MacroH2A.

Suitable for use as control antibody for MacroH2A1 siRNA (h): sc-91790, MacroH2A1 siRNA (m): sc-149209, MacroH2A1 shRNA Plasmid (h): sc-91790-SH, MacroH2A1 shRNA Plasmid (m): sc-149209-SH, MacroH2A1 shRNA (h) Lentiviral Particles: sc-91790-V and MacroH2A1 shRNA (m) Lentiviral Particles: sc-149209-V.

MacroH2A1 (K-14) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of MacroH2A1: 39 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.