

MICAL1 (P-20): sc-161858

BACKGROUND

MICAL1 (microtubule associated monoxygenase, calponin and LIM domain containing 1), also known as MICAL or NICAL, is a 1,067 amino acid protein that localizes to both the cytoplasm and the cytoskeleton and contains one LIM zinc-binding domain and one calponin-homology domain. Expressed in kidney, thymus, spleen, lung and testes, MICAL1 interacts with the SH3 domain of Cas-L and, via this interaction, is thought to function as a cytoskeletal regulator that connects Cas-L to intermediate filaments. MICAL1 also interacts with Rab 1B, plexin-A3 and Vimentin, further regulating cytoskeletal events and possibly playing a role in axonal repulsion. Three isoforms of MICAL1 exist due to alternative splicing events.

REFERENCES

1. Terman, J.R., Mao, T., Pasterkamp, R.J., Yu, H.H. and Kolodkin, A.L. 2002. MICALs, a family of conserved flavoprotein oxidoreductases, function in plexin-mediated axonal repulsion. *Cell* 109: 887-900.
2. Suzuki, T., Nakamoto, T., Ogawa, S., Seo, S., Matsumura, T., Tachibana, K., Morimoto, C. and Hirai, H. 2002. MICAL, a novel CasL interacting molecule, associates with Vimentin. *J. Biol. Chem.* 277: 14933-14941.
3. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 607129. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Weide, T., Teuber, J., Bayer, M. and Barnekow, A. 2003. MICAL1 isoforms, novel rab1 interacting proteins. *Biochem. Biophys. Res. Commun.* 306: 79-86.
5. Bayer, M., Fischer, J., Kremerskothen, J., Ossendorf, E., Matanis, T., Konczal, M., Weide, T. and Barnekow, A. 2005. Identification and characterization of Iporin as a novel interaction partner for rab1. *BMC Cell Biol.* 6: 15.
6. Fischer, J., Weide, T. and Barnekow, A. 2005. The MICAL proteins and rab1: a possible link to the cytoskeleton? *Biochem. Biophys. Res. Commun.* 328: 415-423.
7. Jin, X., Zhang, J., Dai, H., Sun, H., Wang, D., Wu, J. and Shi, Y. 2007. Investigation of the four cooperative unfolding units existing in the MICAL1 CH domain. *Biophys. Chem.* 129: 269-278.
8. Schmidt, E.F., Shim, S.O. and Strittmatter, S.M. 2008. Release of MICAL autoinhibition by semaphorin-plexin signaling promotes interaction with collapsin response mediator protein. *J. Neurosci.* 28: 2287-2297.

CHROMOSOMAL LOCATION

Genetic locus: MICAL1 (human) mapping to 6q21.

SOURCE

MICAL1 (P-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of MICAL1 of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-161858 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

MICAL1 (P-20) is recommended for detection of MICAL1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with MICALCL, MICAL2 or MICAL3.


Suitable for use as control antibody for MICAL1 siRNA (h): sc-95537, MICAL1 shRNA Plasmid (h): sc-95537-SH and MICAL1 shRNA (h) Lentiviral Particles: sc-95537-V.

Molecular Weight of MICAL1: 118 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



MICAL1 (P-20): sc-161858. Western blot analysis of MICAL1 expression in non-transfected 293T: sc-117752 (A), human MICAL1 transfected 293T: sc-116300 (B) and SK-MEL-28 (C) whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.