

# MYBPC1 (T-14): sc-161892

## BACKGROUND

MYBPC1, also known as MYBPCS (myosin-binding protein C, slow-type) or MyBP-C, is a 1,141 amino acid protein that contains three Fibronectin type-III domains and seven Ig-like C2-type domains. Existing as a member of the immunoglobulin superfamily, MYBPC1 functions as a thick filament-associated protein that localizes to striated muscle bands in vertebrae and is thought to modify the activity of select ATPases. Additionally, MYBPC1 may play a role in the modulation of muscle contraction and in the overall structural integrity of the cell. The gene encoding MYBPC1 maps to human chromosome 12, which encodes over 1,100 genes and comprises approximately 4.5% of the human genome. Chromosome 12 is associated with a variety of diseases and afflictions, including hypochondrogenesis, achondrogenesis, Kniest dysplasia, Noonan syndrome and trisomy 12p, which causes facial developmental defects and seizure disorders.

## REFERENCES

- Weber, F.E., et al. 1993. Complete sequence of human fast-type and slow-type muscle myosin-binding-protein C (MyBP-C). Differential expression, conserved domain structure and chromosome assignment. *Eur. J. Biochem.* 216: 661-669.
- Alyonycheva, T.N., et al. 1997. Isoform-specific interaction of the myosin-binding proteins (MyBPs) with skeletal and cardiac myosin is a property of the C-terminal immunoglobulin domain. *J. Biol. Chem.* 272: 20866-20872.
- Welikson, R.E. and Fischman, D.A. 2002. The C-terminal Igl domains of myosin-binding proteins C and H (MyBP-C and MyBPH) are both necessary and sufficient for the intracellular crosslinking of sarcomeric myosin in transfected non-muscle cells. *J. Cell Sci.* 115: 3517-3526.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 160794. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Konno, T., et al. 2003. A novel missense mutation in the myosin binding protein-C gene is responsible for hypertrophic cardiomyopathy with left ventricular dysfunction and dilation in elderly patients. *J. Am. Coll. Cardiol.* 41: 781-786.
- Dhoot, G.K. and Perry, S.V. 2005. Expression of slow skeletal myosin binding C-protein in normal adult mammalian heart. *J. Muscle Res. Cell Motil.* 26: 143-148.
- McGrath, M.J., et al. 2006. Four and a half LIM protein 1 binds myosin-binding protein C and regulates myosin filament formation and sarcomere assembly. *J. Biol. Chem.* 281: 7666-7683.

## CHROMOSOMAL LOCATION

Genetic locus: MYBPC1 (human) mapping to 12q23.2; Mybpc1 (mouse) mapping to 10 C1.

## SOURCE

MYBPC1 (T-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of MYBPC1 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-161892 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

MYBPC1 (T-14) is recommended for detection of MYBPC1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with MYBPC2 or MYBPC3.

MYBPC1 (T-14) is also recommended for detection of MYBPC1 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for MYBPC1 siRNA (h): sc-96079, MYBPC1 siRNA (m): sc-149730, MYBPC1 shRNA Plasmid (h): sc-96079-SH, MYBPC1 shRNA Plasmid (m): sc-149730-SH, MYBPC1 shRNA (h) Lentiviral Particles: sc-96079-V and MYBPC1 shRNA (m) Lentiviral Particles: sc-149730-V.

Molecular Weight of MYBPC1: 128 kDa.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.



Try **MYBPC1 (E-6): sc-515407**, our highly recommended monoclonal alternative to MYBPC1 (T-14).