

OAT1 (N-13): sc-161978

BACKGROUND

The organic anion transporter (OAT) family of proteins mediate the secretion of exogenous and endogenous metabolites from tissues throughout the body. OAT1 (organic anion transporter 1), a 563 amino acid protein, and OAT3 (organic anion transporter 3), a 542 amino acid protein, are two members of the OAT family and are highly expressed in kidneys. Localized specifically to the basolateral membrane, OAT1 and OAT3 are involved in the elimination of toxic organic anions, such as benzylpenicillin and cimetidine, from proximal renal tubules. Via their ability to remove anions from renal tissues, OAT1 and OAT3 are able to regulate the amount of toxins within the kidneys. Additionally, OAT1 functions as an organic anion exchanger that couples the uptake of one organic anion molecule with the efflux of one endogenous dicarboxylic acid molecule, such as ketoglutarate. Four isoforms of OAT1 and three isoforms of OAT3 are expressed due to alternative splicing events.

CHROMOSOMAL LOCATION

Genetic locus: SLC22A6 (human) mapping to 11q12.3.

SOURCE

OAT1 (N-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an N-terminal extracellular domain of OAT1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-161978 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

OAT1 (N-13) is recommended for detection of OAT1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other OAT or OATP family members.

OAT1 (N-13) is also recommended for detection of OAT1 in additional species, including canine.

Suitable for use as control antibody for OAT1 siRNA (h): sc-96343, OAT1 shRNA Plasmid (h): sc-96343-SH and OAT1 shRNA (h) Lentiviral Particles: sc-96343-V.


Molecular Weight (predicted) of OAT1: 62 kDa.

Molecular Weight (observed) of OAT1: 78 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



OAT1 (N-13): sc-161978. Western blot analysis of OAT1 expression in mouse brain tissue extract.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **OAT1 (1F2): sc-293323**, our highly recommended monoclonal alternative to OAT1 (N-13).