

PI 4-kinase II β (S-13): sc-162015

BACKGROUND

Phosphatidylinositol (PI) kinases participate in the first step in activating important cellular effectors, such as PIP2 (phosphatidylinositol bisphosphate) and PTEN. Unlike other PI-kinases, PI 4-kinase family members only phosphorylate phosphatidylinositols, are potently inhibited by adenosine and lack a trans-membrane domain. Total PI 4-kinase activity is dependent upon PI 4-kinase β , PI 4-kinase α , PI 4-kinase II α and PI 4-kinase II β activities. PI 4-kinase II β (phosphatidylinositol 4-kinase type II- β) is a 481 amino acid protein that is recruited to the membrane from the cytosol to stimulate synthesis of PIP2 and, in this way, PI 4-kinase II β contributes to the production of IP3. PI 4-kinase II β may also be involved in the regulation of vesicular trafficking due to its associations with the endoplasmic reticulum and the golgi apparatus. T-cells specific for PI 4-kinase II β lyse normal and malignant hematopoietic cells, suggesting that PI 4-kinase II β may play a role in antitumor immunity.

REFERENCES

- Barylko, B., et al. 2001. A novel family of phosphatidylinositol 4-kinases conserved from yeast to humans. *J. Biol. Chem.* 276: 7705-7708.
- Minogue, S., et al. 2001. Cloning of a human type II phosphatidylinositol 4-kinase reveals a novel lipid kinase family. *J. Biol. Chem.* 276: 16635-16640.
- Balla, A., et al. 2002. Characterization of type II phosphatidylinositol 4-kinase isoforms reveals association of the enzymes with endosomal vesicular compartments. *J. Biol. Chem.* 277: 20041-20050.
- Barylko, B., et al. 2002. Analysis of the catalytic domain of phosphatidylinositol 4-kinase type II. *J. Biol. Chem.* 277: 44366-44375.
- Wei, Y.J., et al. 2002. Type II phosphatidylinositol 4-kinase β is a cytosolic and peripheral membrane protein that is recruited to the plasma membrane and activated by Rac-GTP. *J. Biol. Chem.* 277: 46586-46593.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 612101. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Griffioen, M., et al. 2008. Identification of phosphatidylinositol 4-kinase type II β as HLA class II-restricted target in graft versus leukemia reactivity. *Proc. Natl. Acad. Sci. USA* 105: 3837-3842.

CHROMOSOMAL LOCATION

Genetic locus: PI4K2B (human) mapping to 4p15.2; Pi4k2b (mouse) mapping to 5 C1.

SOURCE

PI 4-kinase II β (S-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of PI 4-kinase II β of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-162015 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

PI 4-kinase II β (S-13) is recommended for detection of PI 4-kinase II β of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive PI 4-kinase II α .

PI 4-kinase II β (S-13) is also recommended for detection of PI 4-kinase II β in additional species, including bovine, porcine and avian.

Suitable for use as control antibody for PI 4-kinase II β siRNA (h): sc-89203, PI 4-kinase II β siRNA (m): sc-152244, PI 4-kinase II β shRNA Plasmid (h): sc-89203-SH, PI 4-kinase II β shRNA Plasmid (m): sc-152244-SH, PI 4-kinase II β shRNA (h) Lentiviral Particles: sc-89203-V and PI 4-kinase II β shRNA (m) Lentiviral Particles: sc-152244-V.

Molecular Weight of PI 4-kinase II β : 54 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.