

Ribosomal Protein S9 (N-15): sc-162106

BACKGROUND

Ribosomes, the organelles that catalyze protein synthesis, are composed of a small subunit (40S) and a large subunit (60S) that consist of over 80 distinct ribosomal proteins. Mammalian ribosomal proteins are encoded by multigene families that contain processed pseudogenes and one functional intron-containing gene within their coding regions. Ribosomal Protein S9 (RPS9), also known as 40S ribosomal protein S9, is a 194 amino acid protein that contains one S4 RNA-binding domain and belongs to the ribosomal protein S4P family. Ribosomal Protein S9 localizes to cytoplasm and has been identified as part of an mRNP granule complex. Like most ribosomal proteins, Ribosomal Protein S9 exists as multiple processed pseudogenes that are scattered throughout the genome. The gene encoding Ribosomal Protein S9 maps to human chromosome 19q13.42.

CHROMOSOMAL LOCATION

Genetic locus: RPS9 (human) mapping to 19q13.42; Rps9 (mouse) mapping to 7 A1.

SOURCE

Ribosomal Protein S9 (N-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of Ribosomal Protein S9 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-162106 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Ribosomal Protein S9 (N-15) is recommended for detection of Ribosomal Protein S9 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other Ribosomal Proteins.

Ribosomal Protein S9 (N-15) is also recommended for detection of Ribosomal Protein S9 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for Ribosomal Protein S9 siRNA (h): sc-97681, Ribosomal Protein S9 siRNA (m): sc-152954, Ribosomal Protein S9 shRNA Plasmid (h): sc-97681-SH, Ribosomal Protein S9 shRNA Plasmid (m): sc-152954-SH, Ribosomal Protein S9 shRNA (h) Lentiviral Particles: sc-97681-V and Ribosomal Protein S9 shRNA (m) Lentiviral Particles: sc-152954-V.


Molecular Weight of Ribosomal Protein S9: 23 kDa.

Positive Controls: Ramos cell lysate: sc-2216, K-562 whole cell lysate: sc-2203 or Ramos nuclear extract: sc-2153.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Ribosomal Protein S9 (N-15): sc-162106. Western blot analysis of Ribosomal Protein S9 expression in Ramos (A), K-562 (B), Raji (C) and SW480 (D) whole cell lysates and A549 (E) and Ramos (F) nuclear extracts.



Ribosomal Protein S9 (N-15): sc-162106. Western blot analysis of Ribosomal Protein S9 expression in 293T (A), MIA PaCa-2 (B) and COLO 320DM (C) whole cell lysates.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **Ribosomal Protein S9 (E-3): sc-390614**, our highly recommended monoclonal alternative to Ribosomal Protein S9 (N-15).