

SCNM1 (C-14): sc-162153

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. SCNM1 (sodium channel modifier 1) is a 230 amino acid protein that contains one matrin-type zinc finger. Localized to the nucleus, SCNM1 is thought to function as an RNA splicing factor that may modify the expression of sodium channel-related proteins. SCNM1 exists as two alternatively spliced isoforms that are encoded by a gene which maps to chromosome 1. Chromosome 1 is the largest human chromosome, spanning about 260 million base pairs and making up 8% of the human genome. Several disorders, including Stickler syndrome, Parkinsons disease, Gaucher disease, malignant melanoma and Usher syndrome, are caused by defects in genes that localize to chromosome 1.

REFERENCES

1. Sprunger, L.K., et al. 1999. Dystonia associated with mutation of the neuronal sodium channel Scn8a and identification of the modifier locus Scnm1 on mouse chromosome 3. *Hum. Mol. Genet.* 8: 471-479.
2. Online Mendelian Inheritance in Man, OMIM[™]. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 608095. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
3. Buchner, D.A., et al. 2003. High-resolution mapping of the sodium channel modifier Scnm1 on mouse chromosome 3 and identification of a 1.3-kb recombination hot spot. *Genomics* 82: 452-459.
4. Buchner, D.A., et al. 2003. SCNM1, a putative RNA splicing factor that modifies disease severity in mice. *Science* 301: 967-969.
5. Howell, V.M., et al. 2007. Evidence for a direct role of the disease modifier SCNM1 in splicing. *Hum. Mol. Genet.* 16: 2506-2516.
6. Howell, V.M., et al. 2008. A targeted deleterious allele of the splicing factor SCNM1 in the mouse. *Genetics* 180: 1419-1427.

CHROMOSOMAL LOCATION

Genetic locus: SCNM1 (human) mapping to 1q21.3; Scnm1 (mouse) mapping to 3 F2.1.

SOURCE

SCNM1 (C-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of SCNM1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-162153 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

SCNM1 (C-14) is recommended for detection of SCNM1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

SCNM1 (C-14) is also recommended for detection of SCNM1 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for SCNM1 siRNA (h): sc-88500, SCNM1 siRNA (m): sc-153263, SCNM1 shRNA Plasmid (h): sc-88500-SH, SCNM1 shRNA Plasmid (m): sc-153263-SH, SCNM1 shRNA (h) Lentiviral Particles: sc-88500-V and SCNM1 shRNA (m) Lentiviral Particles: sc-153263-V.


Molecular Weight of SCNM1: 30 kDa.

Positive Controls: NIH/3T3 nuclear extract: sc-2138, HeLa nuclear extract: sc-2120 or CCRF-CEM nuclear extract: sc-2146.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

DATA



SCNM1 (C-14): sc-162153. Western blot analysis of SCNM1 expression in HeLa (A) and CCRF-CEM (B) nuclear extracts.



SCNM1 (C-14): sc-162153. Western blot analysis of SCNM1 expression in NIH/3T3 nuclear extract.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **SCNM1 (A-6): sc-376328** or **SCNM1 (F-9): sc-376457**, our highly recommended monoclonal alternatives to SCNM1 (C-14).