

SGLT-4 (G-14): sc-162182

BACKGROUND

Glucose is the main source of energy for mammalian cells and its entry is mediated by various transporters. This process involves seven facilitative (GLUT-1 to -7) and multiple members of the SGLT family. SGLT family members use the electrochemical gradient of two sodium ions to transport one glucose molecule. The mRNA of SGLTs increase steadily from the fetal period to maturity along with an increase in their functional activity. SGLT-4 (sodium/glucose cotransporter 4), also known as SLC5A9 (solute carrier family 5 member 9), is a 681 amino acid multi-pass membrane protein that participates in the sodium-dependent transport of D-fructose, D-mannose and D-glucose. A member of the sodium:solute symporter (SSF) family, SGLT-4 is found in small intestine and kidney.

REFERENCES

1. Turk, E., Klisak, I., Bacallao, R., Sparkes, R.S. and Wright, E.M. 1993. Assignment of the human Na⁺/glucose cotransporter gene SGLT1 to chromosome 22q13.1. *Genomics* 17: 752-754.
2. Stümpel, F., Burcelin, R., Jungermann, K. and Thorens, B. 2001. Normal kinetics of intestinal glucose absorption in the absence of GLUT2: evidence for a transport pathway requiring glucose phosphorylation and transfer into the endoplasmic reticulum. *Proc. Natl. Acad. Sci. USA* 98: 11330-11335.
3. Wallner, E.I., Wada, J., Tramonti, G., Lin, S. and Kanwar, Y.S. 2001. Status of glucose transporters in the mammalian kidney and renal development. *Ren. Fail.* 23: 301-310.
4. Wright, E.M., Martín, M.G. and Turk, E. 2003. Intestinal absorption in health and disease—sugars. *Best Pract. Res. Clin. Gastroenterol.* 17: 943-956.
5. Scheepers, A., Joost, H.G. and Schürmann, A. 2004. The glucose transporter families SGLT and GLUT: molecular basis of normal and aberrant function. *JPEN J. Parenter. Enteral. Nutr.* 28: 364-371.
6. Tazawa, S., Yamato, T., Fujikura, H., Hiratochi, M., Itoh, F., Tomae, M., Takemura, Y., Maruyama, H., Sugiyama, T., Wakamatsu, A., Isogai, T. and Isaji, M. 2005. SLC5A9/SGLT4, a new Na⁺-dependent glucose transporter, is an essential transporter for mannose, 1,5-anhydro-D-glucitol, and fructose. *Life Sci.* 76: 1039-1050.

CHROMOSOMAL LOCATION

Genetic locus: SLC5A9 (human) mapping to 1p33; Slc5a9 (mouse) mapping to 4 D1.

SOURCE

SGLT-4 (G-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within a C-terminal cytoplasmic domain of SGLT-4 of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-162182 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

SGLT-4 (G-14) is recommended for detection of SGLT-4 of mouse and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other SGLT family members.

SGLT-4 (G-14) is also recommended for detection of SGLT-4 in additional species, including equine and bovine.

Suitable for use as control antibody for SGLT-4 siRNA (h): sc-88339, SGLT-4 siRNA (m): sc-153420, SGLT-4 shRNA Plasmid (h): sc-88339-SH, SGLT-4 shRNA Plasmid (m): sc-153420-SH, SGLT-4 shRNA (h) Lentiviral Particles: sc-88339-V and SGLT-4 shRNA (m) Lentiviral Particles: sc-153420-V.

Molecular Weight of SGLT-4: 74 kDa.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.