

TMLH (S-12): sc-162336

BACKGROUND

Carnitine is a quaternary ammonium compound that is required for fatty acid transport into the mitochondria. This step is necessary to utilize fatty acids in β -oxidation to obtain usable energy for the citric acid cycle. TMLH (trimethyllysine hydroxylase), also known as ϵ -trimethyllysine 2-oxoglutarate dioxygenase and TML- α -ketoglutarate dioxygenase, is a 421 amino acid mitochondrial matrix protein that converts trimethyllysine (TML) into hydroxytrimethyllysine (HTML), the first of 4 steps in carnitine biosynthesis. Expressed in both fetal and adult human tissue, there are two isoforms of TMLH, designated TMLHa and TMLHb, that are produced as a result of alternative splicing events. Interestingly, TMLHb negatively affect TMLH activity, suggesting that it may act as a crucial physiological negative regulator of TMLH.

REFERENCES

- Vaz, F.M., et al. 2001. Molecular and biochemical characterization of rat ϵ -N-trimethyllysine hydroxylase, the first enzyme of carnitine biosynthesis. *J. Biol. Chem.* 276: 33512-33517.
- Swiegers, J.H., et al. 2002. Carnitine biosynthesis in *Neurospora crassa*: identification of a cDNA coding for epsilon-N-trimethyllysine hydroxylase and its functional expression in *Saccharomyces cerevisiae*. *FEMS Microbiol. Lett.* 210: 19-23.
- Monfregola, J., et al. 2005. Functional analysis of TMLH variants and definition of domains required for catalytic activity and mitochondrial targeting. *J. Cell. Physiol.* 204: 839-847.
- Davis, A.T. and Monroe, T.J. 2005. Carnitine deficiency and supplementation do not affect the gene expression of carnitine biosynthetic enzymes in rats. *J. Nutr.* 135: 761-764.
- Monfregola, J., et al. 2007. Functional characterization of the TMLH gene: promoter analysis, *in situ* hybridization, identification and mapping of alternative splicing variants. *Gene* 395: 86-97.
- Choudhary, C., et al. 2009. Lysine acetylation targets protein complexes and co-regulates major cellular functions. *Science* 325: 834-840.
- Online Mendelian Inheritance in Man, OMIM™. 2009. Johns Hopkins University, Baltimore, MD. MIM Number: 300777. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: TMLHE (human) mapping to Xq28; Tmlhe (mouse) mapping to X.

SOURCE

TMLH (S-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of TMLH of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-162336 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

TMLH (S-12) is recommended for detection of TMLH of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

TMLH (S-12) is also recommended for detection of TMLH in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for TMLH siRNA (h): sc-91155, TMLH shRNA Plasmid (h): sc-91155-SH and TMLH shRNA (h) Lentiviral Particles: sc-91155-V.

Molecular Weight of TMLH precursor: 48 kDa.


Molecular Weight of mature TMLH: 43 kDa.

Positive Controls: JAR cell lysate: sc-2276 or JEG-3 whole cell lysate: sc-364255.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



TMLH (S-12): sc-162336. Western blot analysis of TMLH expression in JAR (A) and JEG-3 (B) whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.