# YIPF2 (M-12): sc-162416



The Power to Question

## **BACKGROUND**

The YIP1 family consists of a group of small membrane proteins that bind Rab GTPases and function in membrane trafficking and vesicle biogenesis. YIPF2 (YIP1 family member 2), also known as FinGER2, is a 316 amino acid multipass membrane protein belonging to the YIP1 family. YIPF2 is encoded by a gene located on human chromosome 19, which consists of over 63 million bases, houses approximately 1,400 genes and is recognized for having the greatest gene density of the human chromosomes. It is the genetic home for a number of immunoglobulin (Ig) superfamily members, including the killer cell and leukocyte Ig-like receptors, a number of ICAMs, the CEACAM and PSG family and Fc receptors (FcRs).

# **REFERENCES**

- 1. Olsen, A., et al. 1994. Gene organization of the pregnancy-specific glycoprotein region on human chromosome 19: assembly and analysis of a 700-kb cosmid contig spanning the region. Genomics 23: 659-668.
- 2. Teglund, S., et al. 1994. The pregnancy-specific glycoprotein (PSG) gene cluster on human chromosome 19: fine structure of the 11 PSG genes and identification of 6 new genes forming a third subgroup within the carcino-embryonic antigen (CEA) family. Genomics 23: 669-684.
- Wang, L., et al. 2000. CEACAM1, a candidate tumor suppressor gene, is abnormally expressed in primary lung cancers. Clin. Cancer Res. 6: 2988-2993.
- 4. Trowsdale, J., et al. 2001. The genomic context of natural killer receptor extended gene families. Immunol. Rev. 181: 20-38.
- 5. Leeb, T. and Müller, M. 2004. Comparative human-mouse-rat sequence analysis of the ICAM gene cluster on HSA 19p13.2 and a 185-kb porcine region from SSC 2q. Gene 343: 239-244.
- Chen, C.Z. and Collins, R.N. 2005. Insights into biological functions across species: examining the role of Rab proteins in YIP1 family function. Biochem. Soc. Trans. 33: 614-618.
- 7. Chen, C.Z. and Collins, R.N. 2005. Analysis and properties of the yeast YIP1 family of Ypt-interacting proteins. Meth. Enzymol. 403: 333-339.
- 8. Barrow, A.D. and Trowsdale, J. 2008. The extended human leukocyte receptor complex: diverse ways of modulating immune responses. Immunol. Rev. 224: 98-123.

# CHROMOSOMAL LOCATION

Genetic locus: Yipf2 (mouse) mapping to 9 A3.

## SOURCE

YIPF2 (M-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of YIPF2 of mouse origin.

## **STORAGE**

Store at 4° C, \*\*D0 NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

#### **PRODUCT**

Each vial contains 200  $\mu g$  IgG in 1.0 ml of PBS with <0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-162416 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## **APPLICATIONS**

YIPF2 (M-12) is recommended for detection of YIPF2 of mouse origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other YIPF family members.

Suitable for use as control antibody for YIPF2 siRNA (m): sc-155405, YIPF2 shRNA Plasmid (m): sc-155405-SH and YIPF2 shRNA (m) Lentiviral Particles: sc-155405-V.

Molecular Weight of YIPF2: 35 kDa.

#### **RECOMMENDED SECONDARY REAGENTS**

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

# **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

### **PROTOCOLS**

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

**Santa Cruz Biotechnology, Inc.** 1.800.457.3801 831.457.3800 fax 831.457.3801 **Europe** +00800 4573 8000 49 6221 4503 0 **www.scbt.com**