

YIPF2 (N-14): sc-162417

BACKGROUND

The YIP1 family consists of a group of small membrane proteins that bind Rab GTPases and function in membrane trafficking and vesicle biogenesis. YIPF2 (YIP1 family member 2), also known as FinGER2, is a 316 amino acid multi-pass membrane protein belonging to the YIP1 family. YIPF2 is encoded by a gene located on human chromosome 19, which consists of over 63 million bases, houses approximately 1,400 genes and is recognized for having the greatest gene density of the human chromosomes. It is the genetic home for a number of immunoglobulin (Ig) superfamily members, including the killer cell and leukocyte Ig-like receptors, a number of ICAMs, the CEACAM and PSG family and Fc receptors (FcRs).

CHROMOSOMAL LOCATION

Genetic locus: YIPF2 (human) mapping to 19p13.2; Yipf2 (mouse) mapping to 9 A3.

SOURCE

YIPF2 (N-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an N-terminal cytoplasmic domain of YIPF2 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-162417 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

YIPF2 (N-14) is recommended for detection of YIPF2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other YIPF family members.

YIPF2 (N-14) is also recommended for detection of YIPF2 in additional species, including equine.

Suitable for use as control antibody for YIPF2 siRNA (h): sc-97660, YIPF2 siRNA (m): sc-155405, YIPF2 shRNA Plasmid (h): sc-97660-SH, YIPF2 shRNA Plasmid (m): sc-155405-SH, YIPF2 shRNA (h) Lentiviral Particles: sc-97660-V and YIPF2 shRNA (m) Lentiviral Particles: sc-155405-V.


Molecular Weight of YIPF2: 35 kDa.

Positive Controls: LNCaP cell lysate: sc-2231, Caki-1 cell lysate: sc-2224 or Saos-2 cell lysate: sc-2235.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



YIPF2 (N-14): sc-162417. Western blot analysis of YIPF2 expression in C4 (A), LNCaP (B), Caki-1 (C) and Saos-2 (D) whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **YIPF2 (A-10): sc-398530**, our highly recommended monoclonal alternative to YIPF2 (N-14).