

zygin 2 (N-16): sc-162442

BACKGROUND

Zygin 2, also known as FEZ2 (fasciculation and elongation protein ζ 2), is a 353 amino acid cytoplasmic protein that belongs to the zygin family. Zygin 2 exists as a difluide-linked homodimer but may form heterodimers with zygin 1. Expressed in non-neural tissues including heart, lung, spleen, muscle, testis, placenta and melanocytes, zygin 2 may be involved in the outgrowth and fasciculation of axons. Existing as two alternatively spliced isoforms, the gene encoding zygin 2 maps to human chromosome 2p22.2 and mouse chromosome 17 E3. The genes encoding zygin 1 and 2 are evolutionarily conserved orthologs to the *C. elegans* gene, UNC-76. Zygin 2 is able to restore partial locomotion and axonal fasciculation to *C. elegans* UNC-76 mutants, which suggest a functional role in axonal outgrowth.

REFERENCES

- Bloom, L. and Horvitz, H.R. 1997. The *Caenorhabditis elegans* gene *unc-76* and its human homologs define a new gene family involved in axonal outgrowth and fasciculation. *Proc. Natl. Acad. Sci. USA* 94: 3414-3419.
- Fujita, T., Ikuta, J., Hamada, J., Okajima, T., Tatematsu, K., Tanizawa, K. and Kuroda, S. 2004. Identification of a tissue-non-specific homologue of axonal fasciculation and elongation protein ζ -1. *Biochem. Biophys. Res. Commun.* 313: 738-744.
- Assmann, E.M., Alborghetti, M.R., Camargo, M.E. and Kobarg, J. 2006. FEZ1 dimerization and interaction with transcription regulatory proteins involves its coiled-coil region. *J. Biol. Chem.* 281: 9869-9881.
- Alborghetti, M.R., Furlan, A.S., Silva, J.C., Paes Leme, A.F., Torriani, I.C. and Kobarg, J. 2010. Human FEZ1 protein forms a disulfide bond mediated dimer: implications for cargo transport. *J. Proteome Res.* 9: 4595-4603.
- Alborghetti, M.R., Furlan, A.S. and Kobarg, J. 2011. FEZ2 has acquired additional protein interaction partners relative to FEZ1: functional and evolutionary implications. *PLoS ONE* 6: e17426.

CHROMOSOMAL LOCATION

Genetic locus: FEZ2 (human) mapping to 2p22.2; Fez2 (mouse) mapping to 17 E3.

SOURCE

zygin 2 (N-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of zygin 2 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-162442 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

zygin 2 (N-16) is recommended for detection of zygin 2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with zygin 1.

zygin 2 (N-16) is also recommended for detection of zygin 2 in additional species, including canine, bovine and porcine.

Suitable for use as control antibody for zygin 2 siRNA (h): sc-94938, zygin 2 siRNA (m): sc-155855, zygin 2 shRNA Plasmid (h): sc-94938-SH, zygin 2 shRNA Plasmid (m): sc-155855-SH, zygin 2 shRNA (h) Lentiviral Particles: sc-94938-V and zygin 2 shRNA (m) Lentiviral Particles: sc-155855-V.


Molecular Weight of zygin 2: 40 kDa.

Positive Controls: zygin 2 (h): 293T Lysate: sc-112470.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



zygin 2 (N-16): sc-162442. Western blot analysis of zygin 2 expression in non-transfected: sc-117752 (A) and human zygin 2 transfected: sc-112470 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.