

AASS (G-13): sc-162469

BACKGROUND

α -aminoadipic semialdehyde synthase (AASS), also designated lysine ketoglutarate reductase (LKR) or saccharopine dehydrogenase (SDH), is a 926 amino acid protein that exists as a homodimer in the mitochondria. AASS acts as a bifunctional enzyme containing the lysine α -ketoglutarate reductase (LKR) and saccharopine dehydrogenase activities that catalyzes the first two steps in lysine degradation. It is widely expressed with highest expression in liver and transcription of the AASS gene is induced upon starvation. Mutations in the gene encoding AASS result in various forms familial hyperlysinemias (FH), autosomal recessive disorders characterized by hyperlysinemia, lysinuria, and variable saccharopinuria. However, no adverse mental or physical effects have been in patients with hyperlysinemia.

CHROMOSOMAL LOCATION

Genetic locus: AASS (human) mapping to 7q31.32; Aass (mouse) mapping to 6 A3.1.

SOURCE

AASS (G-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of AASS of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-162469 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

AASS (G-13) is recommended for detection of AASS of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

AASS (G-13) is also recommended for detection of AASS in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for AASS siRNA (h): sc-89857, AASS siRNA (m): sc-140738, AASS shRNA Plasmid (h): sc-89857-SH, AASS shRNA Plasmid (m): sc-140738-SH, AASS shRNA (h) Lentiviral Particles: sc-89857-V and AASS shRNA (m) Lentiviral Particles: sc-140738-V.

Molecular Weight (predicted) of AASS: 102 kDa.


Molecular Weight (observed) of AASS: 116-128 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, rat liver extract: sc-2395 or NIH/3T3 whole cell lysate: sc-2210.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-2494.

DATA



AASS (G-13): sc-162469. Western blot analysis of AASS expression in HeLa whole cell lysate (A) and rat liver tissue extract (B).



AASS (G-13): sc-162469. Western blot analysis of AASS expression in NIH/3T3 whole cell lysate.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **AASS (H-5): sc-390511** or **AASS (G-7): sc-390536**, our highly recommended monoclonal alternatives to AASS (G-13).