

ARS2 (C-14): sc-162539

BACKGROUND

ARS2 (arsenate (or arsenite-) resistance protein 2), also known as ASR2, is an 876 amino acid protein that belongs to the ARS2 family. Expressed ubiquitously in mammals and localized to the nucleus, ARS2 is evolutionarily conserved (at least 98% sequence identity among mammals) and appears to be essential for early mammalian development with a likely role in vital cellular processes. Mouse embryos lacking ARS2 exhibit excessive apoptosis and die around the time of implantation. In humans, ARS2 is known to interact with RNPS1, a protein involved in the activation of pre-mRNA splicing. In addition, the gene encoding ARS2 is located on chromosome 7 within the region that is commonly deleted in myeloid leukemia. This suggests a possible role of ARS2 in the development of myeloid leukemia. Due to alternative splicing events, ARS2 exists in four isoforms, namely isoform A, isoform B, isoform 3 and isoform 4.

REFERENCES

- Rossmann, T.G. and Wang, Z. 1999. Expression cloning for arsenite-resistance resulted in isolation of tumor-suppressor *fauc* cDNA: possible involvement of the ubiquitin system in arsenic carcinogenesis. *Carcinogenesis* 20: 311-316.
- Rossmann, T.G., Visalli, M.A. and Komissarova, E.V. 2003. *fauc* and its ubiquitin-like domain (FUBI) transforms human osteogenic sarcoma (HOS) cells to anchorage-independence. *Oncogene* 22: 1817-1821.
- Beausoleil, S.A., Jedrychowski, M., Schwartz, D., Elias, J.E., Villen, J., Li, J., Cohn, M.A., Cantley, L.C. and Gygi, S.P. 2004. Large-scale characterization of HeLa cell nuclear phosphoproteins. *Proc. Natl. Acad. Sci. USA* 101: 12130-12135.
- Ordóñez, E., Letek, M., Valbuena, N., Gil, J.A. and Mateos, L.M. 2005. Analysis of genes involved in arsenic resistance in *Corynebacterium glutamicum* ATCC 13032. *Appl. Environ. Microbiol.* 71: 6206-6215.
- Bennetts, J.S., Fowles, L.F., Berkman, J.L., van Bueren, K.L., Richman, J.M., Simpson, F. and Wicking, C. 2006. Evolutionary conservation and murine embryonic expression of the gene encoding the SERTA domain-containing protein CDCA4 (HEPP). *Gene* 374: 153-165.

CHROMOSOMAL LOCATION

Genetic locus: SRRT (human) mapping to 7q22.1; *Srrt* (mouse) mapping to 5 G2.

SOURCE

ARS2 (C-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of ARS2 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-162539 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

ARS2 (C-14) is recommended for detection of ARS2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with ARSA.

ARS2 (C-14) is also recommended for detection of ARS2 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for ARS2 siRNA (h): sc-89387, ARS2 siRNA (m): sc-141277, ARS2 shRNA Plasmid (h): sc-89387-SH, ARS2 shRNA Plasmid (m): sc-141277-SH, ARS2 shRNA (h) Lentiviral Particles: sc-89387-V and ARS2 shRNA (m) Lentiviral Particles: sc-141277-V.

Molecular Weight (predicted) of ARS2: 100 kDa.


Molecular Weight (observed) of ARS2: 131 kDa.

Positive Controls: ARS2 (m): 293T Lysate: sc-118573 or U-251-MG whole cell lysate: sc-364176.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



ARS2 (C-14): sc-162539. Western blot analysis of ARS2 expression in non-transfected 293T: sc-117752 (A), mouse ARS2 transfected 293T: sc-118573 (B) and U-251-MG (C) whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.