

# C14orf104 (S-14): sc-162586

## BACKGROUND

Chromosome 14 contains about 700 genes and 106 million base pairs and makes up about 3.5% of human cellular DNA. Chromosome 14 encodes the presenilin 1 (PSEN1) gene, which is one of the three key genes associated with the development of Alzheimer's disease. The SERPINA1 gene is located on chromosome 14 and when defective leads to the genetic disorder  $\alpha$ 1-antitrypsin deficiency. This disorder is characterized by severe lung complications and liver dysfunction. Notably, the immunoglobulin heavy chain locus is found on chromosome 14 and has been identified as a fusion with the chromosome 19 encoded protein Bcl-3 in the (14;19) translocations found in a variety of B cell malignancies. The C14orf104 gene product has been provisionally designated C14orf104 pending further characterization.

## REFERENCES

1. Heilig, R., et al. 2003. The DNA sequence and analysis of human chromosome 14. *Nature* 421: 601-607.
2. Godbolt, A.K., et al. 2004. A presenilin 1 R278I mutation presenting with language impairment. *Neurology* 63: 1702-1704.
3. Stolk, J., et al. 2006.  $\alpha$ 1-antitrypsin deficiency: current perspective on research, diagnosis, and management. *Int. J. Chron. Obstruct. Pulmon. Dis.* 1: 151-160.
4. Vetrivel, K.S., et al. 2006. Pathological and physiological functions of presenilins. *Mol. Neurodegener.* 1: 4.
5. Albani, D., et al. 2007. Presenilin-1 mutation E318G and familial Alzheimer's disease in the Italian population. *Neurobiol. Aging* 28: 1682-1688.
6. Cruz, P.E., et al. 2007. The promise of gene therapy for the treatment of  $\alpha$ 1 antitrypsin deficiency. *Pharmacogenomics* 8: 1191-1198.
7. Filley, C.M., et al. 2007. The genetics of very early onset Alzheimer disease. *Cogn. Behav. Neurol.* 20: 149-156.
8. Martín-Subero, J.I., et al. 2007. A comprehensive genetic and histopathologic analysis identifies two subgroups of B-cell malignancies carrying a t(14;19)(q32;q13) or variant Bcl-3 translocation. *Leukemia* 21: 1532-1544.

## CHROMOSOMAL LOCATION

Genetic locus: C14orf104 (human) mapping to 14q22.1; 1110034A24Rik (mouse) mapping to 12 C2.

## SOURCE

C14orf104 (S-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of C14orf104 of human origin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-162586 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

C14orf104 (S-14) is recommended for detection of C14orf104 of human origin and 1110034A24Rik of mouse origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other C14orf family members.


Suitable for use as control antibody for C14orf104 siRNA (h): sc-92288, 1110034A24Rik siRNA (m): sc-108178, C14orf104 shRNA Plasmid (h): sc-92288-SH, 1110034A24Rik shRNA Plasmid (m): sc-108178-SH, C14orf104 shRNA (h) Lentiviral Particles: sc-92288-V and 1110034A24Rik shRNA (m) Lentiviral Particles: sc-108178-V.

Molecular Weight of C14orf104 isoform 1: 91 kDa.

Molecular Weight of C14orf104 isoform 2: 86 kDa.

Positive Controls: HCT 116 whole cell lysate.

## DATA



C14orf104 (S-14): sc-162586. Western blot analysis of C14orf104 expression in HCT 116 whole cell lysate.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.