

ERMAP (Q-17): sc-162771

BACKGROUND

ERMAP (erythroblast membrane-associated protein), also known as RD, SC or PRO2801, is a single-pass type one membrane protein that belongs to the immunoglobulin superfamily. Expressed in cord blood, fetal liver and adult bone marrow, ERMAP is thought to function as a cell adhesion molecule in erythroid cells and is responsible for expression of the Scianna/Radin (Sc/Rd) blood group antigen system. The Sc/Rd system is comprised of seven antigens that are present on the surface of red blood cells and have a variety of functions ranging from protein transport to cell adhesion. These seven blood antigens can differ in their expression within a population and may sometimes differ between mother and child. A fetus expressing different blood antigens than its mother may cause the mother to produce antibodies against the fetal blood. This condition is known as hemolytic disease of the newborn (HDN) and is characterized by jaundice, anemia and, in some cases, infant death.

REFERENCES

1. Ye, T.Z., Gordon, C.T., Lai, Y.H., Fujiwara, Y., Peters, L.L., Perkins, A.C. and Chui, D.H. 2000. ERMAP, a gene coding for a novel erythroid specific adhesion/receptor membrane protein. *Gene* 242: 337-345.
2. Su, Y.Y., Gordon, C.T., Ye, T.Z., Perkins, A.C. and Chui, D.H. 2001. Human ERMAP: an erythroid adhesion/receptor transmembrane protein. *Blood Cells Mol. Dis.* 27: 938-949.
3. Xu, H., Foltz, L., Sha, Y., Madlansacay, M.R., Cain, C., Lindemann, G., Vargas, J., Nagy, D., Harriman, B., Mahoney, W. and Schueler, P.A. 2001. Cloning and characterization of human erythroid membrane-associated protein, human ERMAP. *Genomics* 76: 2-4.
4. Wagner, F.F., Poole, J. and Flegel, W.A. 2003. Scianna antigens including Rd are expressed by ERMAP. *Blood* 101: 752-757.
5. Flegel, W.A., Chen, Q., Reid, M.E., Martin, J., Orsini, L.A., Poole, J., Moulds, M.K. and Wagner, F.F. 2005. SCER and SCAN: two novel high-prevalence antigens in the Scianna blood group system. *Transfusion* 45: 1940-1944.
6. Velliquette, R.W. 2005. Review: the Scianna blood group system. *Immunohematology* 21: 70-76.
7. He, X.R., He, Y.Y., Chen, Y. and Ye, T.Z. 2006. Expression of human ERMAP gene in fetal tissues. *Zhongguo Shi Yan Xue Ye Xue Za Zhi* 14: 972-975.

CHROMOSOMAL LOCATION

Genetic locus: ERMAP (human) mapping to 1p34.2; Ermap (mouse) mapping to 4 D2.1.

SOURCE

ERMAP (Q-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within a cytoplasmic domain of ERMAP of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-162771 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

ERMAP (Q-17) is recommended for detection of ERMAP of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

ERMAP (Q-17) is also recommended for detection of ERMAP in additional species, including equine and bovine.

Suitable for use as control antibody for ERMAP siRNA (h): sc-88045, ERMAP siRNA (m): sc-144934, ERMAP shRNA Plasmid (h): sc-88045-SH, ERMAP shRNA Plasmid (m): sc-144934-SH, ERMAP shRNA (h) Lentiviral Particles: sc-88045-V and ERMAP shRNA (m) Lentiviral Particles: sc-144934-V.

Molecular Weight of ERMAP: 53 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **ERMAP (YS-6): sc-81790**, our highly recommended monoclonal alternative to ERMAP (Q-17).