

GLIPR1 (P-14): sc-162876

BACKGROUND

GLIPR1 (GLI pathogenesis-related 1), also known as GLIPR, RTVP1 or CRISP7, is a 266 amino acid single-pass membrane protein belonging to the cysteine-rich secretory protein (CRISP) family. GLIPR1 also shares similarity with the pathogenesis-related protein (PR) superfamily, and may function as a p53 target gene with tumor suppressor functions. While ubiquitously expressed, GLIPR1 is found at highest levels in heart, lung, kidney, placenta, liver, skeletal muscle and cell lines derived from tumors of the nervous system. GLIPR1 expression is induced by p53 overexpression, exposure to γ irradiation and doxorubicin. The gene encoding GLIPR1 maps to human chromosome 12q21.2 and while alternatively spliced isoforms of GLIPR1 are known to exist, not all variants have been characterized.

REFERENCES

1. Murphy, E.V., et al. 1995. The human glioma pathogenesis-related protein is structurally related to plant pathogenesis-related proteins and its gene is expressed specifically in brain tumors. *Gene* 159: 131-135.
2. Rich, T., et al. 1996. RTVP-1, a novel human gene with sequence similarity to genes of diverse species, is expressed in tumor cell lines of glial but not neuronal origin. *Gene* 180: 125-130.
3. Ren, C., et al. 2002. mRTVP-1, a novel p53 target gene with proapoptotic activities. *Mol. Cell. Biol.* 22: 3345-3357.
4. Rosenzweig, T., et al. 2006. Related to testes-specific, vespid, and pathogenesis protein-1 (RTVP-1) is overexpressed in gliomas and regulates the growth, survival, and invasion of glioma cells. *Cancer Res.* 66: 4139-4148.
5. Ren, C., et al. 2006. Identification and characterization of RTVP1/GLIPR1-like genes, a novel p53 target gene cluster. *Genomics* 88: 163-172.
6. Xiang, C., et al. 2007. Cloning and characterization of human RTVP-1b, a novel splice variant of RTVP-1 in glioma cells. *Biochem. Biophys. Res. Commun.* 362: 612-618.

CHROMOSOMAL LOCATION

Genetic locus: GLIPR1 (human) mapping to 12q21.2.

SOURCE

GLIPR1 (P-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of GLIPR1 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-162876 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

GLIPR1 (P-14) is recommended for detection of GLIPR1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

GLIPR1 (P-14) is also recommended for detection of GLIPR1 in additional species, including equine, bovine and porcine.

Suitable for use as control antibody for GLIPR1 siRNA (h): sc-96218, GLIPR1 shRNA Plasmid (h): sc-96218-SH and GLIPR1 shRNA (h) Lentiviral Particles: sc-96218-V.


Molecular Weight of GLIPR1: 30 kDa.

Positive Controls: T98G Cell Lysate: sc-2294.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



GLIPR1 (P-14): sc-162876. Western blot analysis of GLIPR1 expression in T98G whole cell lysate.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **GLIPR1 (8D9): sc-517141**, our highly recommended monoclonal alternative to GLIPR1 (P-14).