

MOBKL2B (G-11): sc-163087

BACKGROUND

MOBKL2B (MOB1, Mps one binder kinase activator-like 2B) is a 216 amino acid protein that is related to the yeast MOB1 protein. Yeast MOB1 binds Mps1, a protein kinase essential for spindle pole body duplication and mitotic checkpoint regulation. Belonging to the MOB1/phocea family, MOBKL2B is suggested to regulate the activity of kinases. The gene encoding MOBKL2B is located on human chromosome 9, which consists of about 145 million bases and 4% of the human genome and encodes nearly 900 genes. Chromosome 9 is considered to play a role in gender determination, deletion of the distal portion of 9p can lead to development of male to female sex reversal, the phenotype of a female with a male X,Y genotype.

REFERENCES

1. van Slegtenhorst, M., de Hoogt, R., Hermans, C., Nellist, M., Janssen, B., Verhoef, S., Lindhout, D., van den Ouweland, A., Halley, D., Young, J., Burley, M., Jeremiah, S., Woodward, K., Nahmias, J., Fox, M., et al. 1997. Identification of the tuberous sclerosis gene TSC1 on chromosome 9q34. *Science* 277: 805-808.
2. Hergovich, A., Schmitz, D. and Hemmings, B.A. 2006. The human tumour suppressor LATS1 is activated by human MOB1 at the membrane. *Biochem. Biophys. Res. Commun.* 345: 50-58.
3. Sasaki, H., Kawano, O., Endo, K., Suzuki, E., Yukiue, H., Kobayashi, Y., Yano, M. and Fujii, Y. 2007. Human MOB1 expression in non-small-cell lung cancer. *Clin. Lung Cancer* 8: 273-276.
4. Praskova, M., Xia, F. and Avruch, J. 2008. MOBKL1A/MOBKL1B phosphorylation by MST1 and MST2 inhibits cell proliferation. *Curr. Biol.* 18: 311-321.
5. Beà, S., Salaverria, I., Armengol, L., Pinyol, M., Fernández, V., Hartmann, E.M., Jares, P., Amador, V., Hernández, L., Navarro, A., Ott, G., Rosenwald, A., Estivill, X. and Campo, E. 2009. Uniparental disomies, homozygous deletions, amplifications, and target genes in mantle cell lymphoma revealed by integrative high-resolution whole-genome profiling. *Blood* 113: 3059-3069.
6. Wilmeth, L.J., Shrestha, S., Montañó, G., Rashe, J. and Shuster, C.B. 2010. Mutual dependence of Mob1 and the chromosomal passenger complex for localization during mitosis. *Mol. Biol. Cell* 21: 380-392.

CHROMOSOMAL LOCATION

Genetic locus: MOBKL2B (human) mapping to 9p21.2; Mobkl2b (mouse) mapping to 4 A5.

SOURCE

MOBKL2B (G-11) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of MOBKL2B of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-163087 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

MOBKL2B (G-11) is recommended for detection of MOBKL2B of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with MOBKL2A and MOBKL2C.

MOBKL2B (G-11) is also recommended for detection of MOBKL2B in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for MOBKL2B siRNA (h): sc-92685, MOBKL2B siRNA (m): sc-149492, MOBKL2B shRNA Plasmid (h): sc-92685-SH, MOBKL2B shRNA Plasmid (m): sc-149492-SH, MOBKL2B shRNA (h) Lentiviral Particles: sc-92685-V and MOBKL2B shRNA (m) Lentiviral Particles: sc-149492-V.

Molecular Weight of MOBKL2B: 25 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.