

PRKRIP1 (D-13): sc-163260

BACKGROUND

PRKRIP1 (PKR interacting protein 1, IL-11 inducible), also known as C114, is a 184 amino acid IL-11 induced protein that is expressed in brain, liver, testis and kidney. Localizing to the nucleus with highest expression in the nucleolus, PRKRIP1 contains an arginine-rich region and displays double-stranded RNA (dsRNA)-binding activity. Via its N-terminus, PRKRIP1 interacts with PKR (protein kinase RNA-activated) and functions to inhibit or negatively regulate PKR activity. The inhibition of PKR leads to the inhibition of eIF2 α phosphorylation. eIF2 α , a component of the eukaryotic initiation complex, functions to bind tRNA to the 40S ribosomal subunit (in a GTP-dependent manner), thereby initiating translation. The phosphorylation state of eIF2 α controls the rate of tRNA translation and, when phosphorylated, translation/protein synthesis is inhibited. This suggests that PRKRIP1, through its ability to inhibit eIF2 α phosphorylation and promote protein synthesis, may play a role in adipogenesis.

REFERENCES

1. Yin, Z., Haynie, J., Williams, B.R. and Yang, Y.C. 2003. C114 is a novel IL-11-inducible nuclear double-stranded RNA-binding protein that inhibits protein kinase R. *J. Biol. Chem.* 278: 22838-22845.
2. Will, C.L., Schneider, C., Hossbach, M., Urlaub, H., Rauhut, R., Elbashir, S., Tuschl, T. and Lührmann, R. 2004. The human 18S U11/U12 snRNP contains a set of novel proteins not found in the U2-dependent spliceosome. *RNA* 10: 929-941
3. Rual, J.F., Venkatesan, K., Hao, T., Hirozane-Kishikawa, T., Dricot, A., Li, N., Berriz, G.F., Gibbons, F.D., Dreze, M., Ayivi-Guedehoussou, N., Klitgord, N., Simon, C., Boxem, M., Milstein, S., Rosenberg, J., Goldberg, D.S., Zhang, L.V., Wong, S.L., Franklin, G., Li, S., Albala, J.S., et al. 2005. Towards a proteome-scale map of the human protein-protein interaction network. *Nature* 437: 1173-1178.
4. Kimura, K., Wakamatsu, A., Suzuki, Y., Ota, T., Nishikawa, T., Yamashita, R., Yamamoto, J., Sekine, M., Tsuritani, K., Wakaguri, H., Ishii, S., Sugiyama, T., Saito, K., Isono, Y., Irie, R., Kushida, N., Yoneyama, T., Otsuka, R., Kanda, K., Yokoi, T., Kondo, H., et al. 2006. Diversification of transcriptional modulation: large-scale identification and characterization of putative alternative promoters of human genes. *Genome Res.* 16: 55-65.

CHROMOSOMAL LOCATION

Genetic locus: PRKRIP1 (human) mapping to 7q22.1; Prkrip1 (mouse) mapping to 5 G2.

SOURCE

PRKRIP1 (D-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of PRKRIP1 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-163260 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

PRKRIP1 (D-13) is recommended for detection of PRKRIP1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

PRKRIP1 (D-13) is also recommended for detection of PRKRIP1 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for PRKRIP1 siRNA (h): sc-89660, PRKRIP1 siRNA (m): sc-152469, PRKRIP1 shRNA Plasmid (h): sc-89660-SH, PRKRIP1 shRNA Plasmid (m): sc-152469-SH, PRKRIP1 shRNA (h) Lentiviral Particles: sc-89660-V and PRKRIP1 shRNA (m) Lentiviral Particles: sc-152469-V.

Molecular Weight of PRKRIP1: 21 kDa.

Positive Controls: HeLa nuclear extract: sc-2120.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.