

SP-100 (N-20): sc-16328

BACKGROUND

The human SP100 gene encodes an autoantigen that co-localizes with PML and NDP52 in distinct nuclear domains, called nuclear dots (NDs) or ND10 nuclear bodies. Papova-, adeno-, and herpesviruses begin their transcription and DNA-replication at NDs, which play a role in autoimmunity, viral infections and in the etiology of acute promyelocytic leukemia. SP-100 is an interferon-inducible protein that has two splice variants. One splice variant contains a highly conserved copy of the DNA-binding high mobility group 1 protein sequence, and thus represents a novel HMG-box protein. This alternatively spliced variant of SP-100 has a unique expression and localization pattern that is distinct from the SP-100 full-length protein. The SP100 protein is covalently modified by the small ubiquitin-related protein SUMO-1. SP-100 contains a functional nuclear localization signal and an ND-targeting region, which overlaps with the SP-100 homodimerization domain. The homodimerization/ND-targeting region is considered a novel protein motif, termed HSR domain. SP-100 is also found to interact with Bright (B cell regulator of IgH transcription), which in lymphoid cells also interacts with LYSP100/SP140, the lymphoid-restricted homolog of SP100 (8).

REFERENCES

- Grotzinger, T., et al. 1996. The interferon (IFN)-stimulated gene SP-100 promoter contains an IFN- γ activation site and an imperfect IFN-stimulated response element which mediate type I IFN inducibility. *J. Biol. Chem.* 271: 25253-25260.
- Weichenhan, D., et al. 1997. Structure and expression of the murine SP-100 nuclear dot gene. *Genomics* 43: 298-306.
- Seeler, J.S., et al. 1998. Interaction of SP-100 with HP1 proteins: a link between the promyelocytic leukemia-associated nuclear bodies and the chromatin compartment. *Proc. Natl. Acad. Sci. USA* 95: 7316-7321.
- Guldner, H.H., et al. 1999. Splice variants of the nuclear dot-associated SP-100 protein contain homologies to HMG 1 and a human nuclear phosphoprotein-box motif. *J. Cell Sci.* 112: 733-747.
- Sternsdorf, T., et al. 1999. The nuclear dot protein SP-100, characterization of domains necessary for dimerization, subcellular localization, and modification by small ubiquitin-like modifiers. *J. Biol. Chem.* 274: 12555-12566.

CHROMOSOMAL LOCATION

Genetic locus: SP100 (human) mapping to 2q37.

SOURCE

SP-100 (N-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of SP-100 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-16328 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

SP-100 (N-20) is recommended for detection of SP-100 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for SP-100 siRNA (h): sc-41032, SP-100 shRNA Plasmid (h): sc-41032-SH and SP-100 shRNA (h) Lentiviral Particles: sc-41032-V.


Molecular Weight of SP-100: 53 kDa.

Positive Controls: HeLa nuclear extract: sc-2120, C32 nuclear extract: sc-2136 or BJAB nuclear extract: sc-2145.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



SP-100 (N-20): sc-16328. Western blot analysis of SP-100 expression in BJAB (A), HeLa (B) and C32 (C) nuclear extracts.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



Try **SP-100 (1G6): sc-293458**, our highly recommended monoclonal alternative to SP-100 (N-20).