# RhCG (E-14): sc-163288



The Power to Questio

#### **BACKGROUND**

RhCG (Rhesus blood group family type C glycoprotein), also known as RHGK (Rh glycoprotein kidney), PDRC2 (tumor-related protein DRC2) or SLC42A3, is a multi-pass membrane protein with three potential N-glycosylation sites that belongs to the Rh subfamily of the ammonium transporter family. Expressed in a wide variety of tissues with predominant expression in adult and fetal kidney, RhCG localizes to the apical and/or basolateral cell membrane (depending on the species) and is believed to function as a non-erythroid, bidirectional, electroneutral ammonium transporter. Specifically, RhCG is found throughout the kidney in sites such as the CNT (connecting segment), the DCT (distal convoluted tubule), the collecting duct and the ICT (initial collecting tubule), that are major contributors to renal ammonia secretion. This suggests that RhCG plays an important role in renal ammonia metabolism.

## **REFERENCES**

- Liu, Z., et al. 2000. Characterization of human RhCG and mouse Rhcg as novel nonerythroid Rh glycoprotein homologues predominantly expressed in kidney and testis. J. Biol. Chem. 275: 25641-25651.
- 2. Heitman, J. and Agre, P. 2000. A new face of the Rhesus antigen. Nat. Genet. 26: 258-259.
- Marini, A.M., et al. 2000. The human Rhesus-associated RhAG protein and a kidney homologue promote ammonium transport in yeast. Nat. Genet. 26: 341-344.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 605381. World Wide Web URL: http://www.ncbi.nlm.nih.gov/omim/
- 5. Khademi, S., et al. 2004. Mechanism of ammonia transport by Amt/MEP/Rh: structure of AmtB at 1.35 A. Science 305: 1587-1594.
- Han, K.H., et al. 2006. Expression of the ammonia transporter, RhC glycoprotein, in normal and neoplastic human kidney. J. Am. Soc. Nephrol. 17: 2670-2679.
- 7. Callebaut, I., et al. 2006. Hydrophobic cluster analysis and modeling of the human Rh protein three-dimensional structures. Transfus. Clin. Biol. 13: 70-84.
- 8. Kim, H.Y., et al. 2007. Effect of reduced renal mass on renal ammonia transporter family, RhC glycoprotein and RhB glycoprotein, expression. Am. J. Physiol. Renal Physiol. 293: F1238-F1247.

# CHROMOSOMAL LOCATION

Genetic locus: Rhcg (mouse) mapping to 7 D3.

## **SOURCE**

RhCG (E-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within a C-terminal cytoplasmic domain of RhCG of mouse origin.

#### **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

#### **PRODUCT**

Each vial contains 200  $\mu g$  lgG in 1.0 ml of PBS with <0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-163288 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

#### **APPLICATIONS**

RhCG (E-14) is recommended for detection of RhCG of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for RhCG siRNA (m): sc-152850, RhCG shRNA Plasmid (m): sc-152850-SH and RhCG shRNA (m) Lentiviral Particles: sc-152850-V.

Molecular Weight of RhCG: 58 kDa.

Positive Controls: PC-12 cell lysate: sc-2250.

#### **RECOMMENDED SECONDARY REAGENTS**

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

#### **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

## **PROTOCOLS**

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

**Santa Cruz Biotechnology, Inc.** 1.800.457.3801 831.457.3801 **Europe** +00800 4573 8000 49 6221 4503 0 **www.scbt.com**