

Sec24D (L-14): sc-163346

BACKGROUND

Sec24D (Sec24-related protein D) is ubiquitously expressed but exhibits predominant expression in heart, placenta, liver and pancreas. Sec24D belongs to the Sec24 subfamily of the Sec23/Sec24 family of proteins. Members of this family participate in vesicle trafficking from the endoplasmic reticulum (ER) to the Golgi apparatus. Sec24D is one of four mammalian proteins, namely Sec24A, Sec24B, Sec24C and Sec24D, that are highly related to the *Saccharomyces cerevisiae* protein Sec24, a component of the coat protein complex COPII that mediates the selective export of membrane proteins from the ER. Similar to its yeast counterpart, Sec24D functions as a component of the cytoplasmic COPII complex. The COPII complex acts as a coat, covering ER-derived transport vesicles and promoting the transport of secretory proteins to the Golgi apparatus. Sec24D shares 50% amino acid identity with the ubiquitously expressed Sec24C protein.

REFERENCES

- Nagase, T., Ishikawa, K., Suyama, M., Kikuno, R., Miyajima, N., Tanaka, A., Kotani, H., Nomura, N. and Ohara, O. 1998. Prediction of the coding sequences of unidentified human genes. XI. The complete sequences of 100 new cDNA clones from brain which code for large proteins *in vitro*. DNA Res. 5: 277-286.
- Tang, B.L., Kausalya, J., Low, D.Y., Lock, M.L. and Hong, W. 1999. A family of mammalian proteins homologous to yeast Sec24p. Biochem. Biophys. Res. Commun. 258: 679-684.
- Online Mendelian Inheritance in Man, OMIM[™]. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 607186. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Farhan, H., Korkhov, V.M., Paulitschke, V., Dorostkar, M.M., Scholze, P., Kudlacek, O., Freissmuth, M. and Sitte, H.H. 2004. Two discontinuous segments in the carboxyl-terminus are required for membrane targeting of the rat γ -aminobutyric acid transporter-1 (GAT1). J. Biol. Chem. 279: 28553-28563.
- Ishizuka, N., Minami, K., Okumachi, A., Okuno, M. and Seino, S. 2007. Induction by NeuroD of the components required for regulated exocytosis. Biochem. Biophys. Res. Commun. 354: 271-277.
- Wendeler, M.W., Paccaud, J.P. and Hauri, H.P. 2007. Role of Sec24 isoforms in selective export of membrane proteins from the endoplasmic reticulum. EMBO Rep. 8: 258-264.
- Farhan, H., Reiterer, V., Korkhov, V.M., Schmid, J.A., Freissmuth, M. and Sitte, H.H. 2007. Concentrative export from the endoplasmic reticulum of the γ -aminobutyric acid transporter 1 requires binding to Sec24D. J. Biol. Chem. 282: 7679-7689.
- Fernández-Sánchez, E., Díez-Guerra, F.J., Cubelos, B., Giménez, C. and Zafra, F. 2008. Mechanisms of endoplasmic-reticulum export of glycine transporter-1 (GLYT1). Biochem. J. 409: 669-681.

CHROMOSOMAL LOCATION

Genetic locus: SEC24D (human) mapping to 4q26; Sec24d (mouse) mapping to 3 G1.

SOURCE

Sec24D (L-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Sec24D of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-163346 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Sec24D (L-14) is recommended for detection of Sec24D of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other Sec24 family members.

Sec24D (L-14) is also recommended for detection of Sec24D in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for Sec24D siRNA (h): sc-89055, Sec24D siRNA (m): sc-153313, Sec24D shRNA Plasmid (h): sc-89055-SH, Sec24D shRNA Plasmid (m): sc-153313-SH, Sec24D shRNA (h) Lentiviral Particles: sc-89055-V and Sec24D shRNA (m) Lentiviral Particles: sc-153313-V.

Molecular Weight of Sec24D: 113 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



MONOS
Satisfaction
Guaranteed

Try **Sec24D (34.1): sc-101268**, our highly recommended monoclonal alternative to Sec24D (L-14).