

Factor VII (K-20): sc-16343

BACKGROUND

Homeostasis following tissue injury involves the deployment of essential plasma procoagulants (prothrombin, and Factors X, IX, V, and VIII), which are involved in a blood coagulation cascade that leads to the formation of insoluble fibrin clots and the promotion of platelet aggregation. Coagulation Factor VII (serum prothrombin conversion accelerator, proconvertin, F7, Factor VII) is a 406 amino acid, vitamin K-dependent, single chain serine protease that is synthesized in the liver and circulates as an inactive precursor. Factor IX A, Factor X A, Factor XII A, or thrombin mediated proteolytic cleavage of Factor VII at Arg152-Ile153 generates Factor VII A, an active serine protease composed of a catalytic heavy chain disulfide linked to a light chain, containing 2 EGF-like domains. Mutations at the F7 locus that lead to Factor VII deficiencies are generally asymptomatic or phenotypically uncharacterized, with hemorrhagic diathesis occurring at extremely low levels.

REFERENCES

1. Davie, E.W. and Fujikawa, K. 1975. Basic mechanisms in blood coagulation. *Annu. Rev. Biochem.* 44: 799-829.
2. Hagen, F.S., et al. 1986. Characterization of a cDNA coding for human Factor VII. *Proc. Natl. Acad. Sci. USA* 83: 2412-2416.
3. O'Hara, P.J., et al. 1987. Nucleotide sequence of the gene coding for human Factor VII, a vitamin K-dependent protein participating in blood coagulation. *Proc. Natl. Acad. Sci. USA* 84: 5158-5162.
4. Davie, E.W., et al. 1991. The coagulation cascade: initiation, maintenance, and regulation. *Biochemistry* 30: 10363-10370.
5. Chambers, R.C., et al. 2000. Thrombin is a potent inducer of connective tissue growth factor production via proteolytic activation of protease-activated receptor-1. *J. Biol. Chem.* 275: 35584-35591.
6. Millar, D.S., et al. 2000. Molecular analysis of the genotype-phenotype relationship in Factor VII deficiency. *Hum. Genet.* 107: 327-342.
7. Online Mendelian Inheritance in Man, OMIM[™]. 2001. Johns Hopkins University, Baltimore, MD. MIM Number: 227500. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
8. McVey, J.H., et al. 2001. Factor VII deficiency and the FVII mutation database. *Hum. Mutat.* 17: 3-17.

CHROMOSOMAL LOCATION

Genetic locus: F7 (human) mapping to 13q34.

SOURCE

Factor VII (K-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Factor VII of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-16343 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Factor VII (K-20) is recommended for detection of Factor VII of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Factor VII siRNA (h): sc-40401, Factor VII shRNA Plasmid (h): sc-40401-SH and Factor VII shRNA (h) Lentiviral Particles: sc-40401-V.

Molecular Weight of Factor VII: 200/80/73/50/43 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941. 3) Immunohistochemistry: use ImmunoCruz[™]: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



Factor VII (K-20): sc-16343. Immunoperoxidase staining of formalin fixed, paraffin-embedded human gall bladder tissue showing cytoplasmic and membrane staining of glandular cells.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **Factor VII (CaFVII-22): sc-101369**, our highly recommended monoclonal alternative to Factor VII (K-20).