

TSCOT (C-13): sc-163477

BACKGROUND

TSCOT (Thymic stromal cotransporter homolog) is a 475 amino acid multi-pass membrane protein that belongs to the SLC46A family. TSCOT is believed to act as a cell membrane transporter and is likely to be found glycosylated. TSCOT is encoded by a gene that maps to human chromosome 9, which consists of about 145 million bases and 4% of the human genome and encodes nearly 900 genes. Considered to play a role in gender determination, deletion of the distal portion of 9p can lead to development of male to female sex reversal, the phenotype of a female with a male X,Y genotype. Hereditary hemorrhagic telangiectasia, which is characterized by harmful vascular defects, is associated with the chromosome 9 gene encoding endoglin protein, ENG. Familial dysautonomia is also associated with chromosome 9 through the gene IKBKAP. Notably, chromosome 9 encompasses the largest interferon family gene cluster.

REFERENCES

- Chen, C., Kim, M.G., Soo Lyu, M., Kozak, C.A., Schwartz, R.H. and Flomerfelt, F.A. 2000. Characterization of the mouse gene, human promoter and human cDNA of TSCOT reveals strong interspecies homology. *Biochim. Biophys. Acta* 1493: 159-169.
- Kim, M.G., Flomerfelt, F.A., Lee, K.N., Chen, C. and Schwartz, R.H. 2000. A putative 12 transmembrane domain cotransporter expressed in thymic cortical epithelial cells. *J. Immunol.* 164: 3185-3192.
- Obermann, H., Wingbermühle, A., Münz, S. and Kirchhoff, C. 2003. A putative 12-transmembrane domain cotransporter associated with apical membranes of the epididymal duct. *J. Androl.* 24: 542-556.
- Yang, S.J., Ahn, S., Park, C.S., Choi, S. and Kim, M.G. 2005. Identifying subpopulations of thymic epithelial cells by flow cytometry using a new specific thymic epithelial marker, Ly110. *J. Immunol. Methods.* 297: 265-270.
- Engelmark, M.T., Ivansson, E.L., Magnusson, J.J., Gustavsson, I.M., Beskow, A.H., Magnusson, P.K. and Gyllensten, U.B. 2006. Identification of susceptibility loci for cervical carcinoma by genome scan of affected sib-pairs. *Hum. Mol. Genet.* 15: 3351-3360.
- Engelmark, M.T., Ivansson, E.L., Magnusson, J.J., Gustavsson, I.M., Wyöni, P.I., Ingman, M., Magnusson, P.K. and Gyllensten, U.B. 2008. Polymorphisms in 9q32 and TSCOT are linked to cervical cancer in affected sib-pairs with high mean age at diagnosis. *Hum. Genet.* 123: 437-443.
- Ahn, S., Lee, G., Yang, S.J., Lee, D., Lee, S., Shin, H.S., Kim, M.C., Lee, K.N., Palmer, D.C., Theoret, M.R., Jenkinson, E.J., Anderson, G., Restifo, N.P. and Kim, M.G. 2008. TSCOT⁺ thymic epithelial cell-mediated sensitive CD4 tolerance by direct presentation. *PLoS Biol.* 6: e191.
- Bateup, H.S., Takasaki, K.T., Saulnier, J.L., Denefrio, C.L. and Sabatini, B.L. 2011. Loss of Tsc1 *in vivo* impairs hippocampal mGluR-LTD and increases excitatory synaptic function. *J. Neurosci.* 31: 8862-8869.

CHROMOSOMAL LOCATION

Genetic locus: SLC46A2 (human) mapping to 9q32.

SOURCE

TSCOT (C-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within a C-terminal cytoplasmic domain of TSCOT of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-163477 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

TSCOT (C-13) is recommended for detection of TSCOT of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for TSCOT siRNA (h): sc-92745, TSCOT shRNA Plasmid (h): sc-92745-SH and TSCOT shRNA (h) Lentiviral Particles: sc-92745-V.

Molecular Weight of TSCOT: 51 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.