C10orf107 (C-14): sc-163761



The Power to Question

BACKGROUND

C10orf107 (chromosome 10 open reading frame 107) is a 208 amino acid protein that is encoded by a gene located on human chromosome 10. Spanning nearly 135 million base pairs, chromosome 10 makes up approximately 4.5% of total DNA in cells and encodes nearly 1,200 genes. Several proteincoding genes, including those that encode for chemokines, cadherins, excision repair proteins, early growth response factors (Egrs) and fibroblast growth receptors (FGFRs), are located on chromosome 10. Defects in some of the genes that map to chromosome 10 are associated with Charcot-Marie Tooth disease, Jackson-Weiss syndrome, Usher syndrome, nonsyndromatic deafness, Wolman's syndrome, Cowden syndrome, multiple endocrine neoplasia type 2 and porphyria.

REFERENCES

- Jabs, E.W., et al. 1994. Jackson-Weiss and Crouzon syndromes are allelic with mutations in fibroblast growth factor receptor 2. Nat. Genet. 8: 275-279.
- Deloukas, P., et al. 2000. Report of the third international workshop on human chromosome 10 mapping and sequencing 1999. Cytogenet. Cell Genet. 90: 1-12.
- 3. Gilbert, F. 2001. Chromosome 10. Genet. Test. 5: 69-82.
- 4. Berger, P., et al. 2002. Molecular cell biology of Charcot-Marie-Tooth disease. Neurogenetics 4: 1-15.
- 5. Nonneman, D., et al. 2004. Comparative mapping of human chromosome 10 to pig chromosomes 10 and 14. Anim. Genet. 35: 338-343.
- Deloukas, P., et al. 2004. The DNA sequence and comparative analysis of human chromosome 10. Nature 429: 375-381.
- 7. Chen, L., et al. 2005. Roles of FGF signaling in skeletal development and human genetic diseases. Front. Biosci. 10: 1961-1976.
- 8. Cho, M.Y., et al. 2008. First report of ovarian dysgerminoma in Cowden syndrome with germline PTEN mutation and PTEN-related 10q loss of tumor heterozygosity. Am. J. Surg. Pathol. 32: 1258-1264.

CHROMOSOMAL LOCATION

Genetic locus: C10orf107 (human) mapping to 10q21.2; 1700040L02Rik (mouse) mapping to 10 B5.2.

SOURCE

C10orf107 (C-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of C10orf107 of human origin.

PRODUCT

Each vial contains 200 μg lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-163761 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

C10orf107 (C-14) is recommended for detection of C10orf107 of human origin and 1700040L02Rik of mouse origin of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other C10orf family members.

Suitable for use as control antibody for C10orf107 siRNA (h): sc-90382, 1700040L02Rik siRNA (m): sc-108435, C10orf107 shRNA Plasmid (h): sc-90382-SH, 1700040L02Rik shRNA Plasmid (m): sc-108435-SH, C10orf107 shRNA (h) Lentiviral Particles: sc-90382-V and 1700040L02Rik shRNA (m) Lentiviral Particles: sc-108435-V.

Molecular Weight of C10orf107: 24 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 Fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com