

FAM188A (N-12): sc-163796

BACKGROUND

FAM188A, also known as Dermal papilla-derived protein 5 and CARP, is a 445 amino acid nuclear protein that likely has pro-apoptotic function. It was originally thought that FAM188A contained a CARD domain, however none of the prediction tools have verified this. FAM188A is highly expressed in kidney, heart and skeletal muscle and is weakly expressed in brain and liver. There are two isoforms of FAM188A that are produced as a result of alternative splicing events. The gene encoding FAM188A maps to human chromosome 10, which contains over 800 genes and 135 million nucleotides. PTEN is an important tumor suppressor gene located on chromosome 10 and, when defective, causes a genetic predisposition to cancer development known as Cowden syndrome. The chromosome 10 encoded gene ERCC6 is important for DNA repair and is linked to Cockayne syndrome which is characterized by extreme photosensitivity and premature aging. Tetrahydrobiopterin deficiency and a number of syndromes involving defective skull and facial bone fusion are also linked to chromosome 10. As with most trisomies, trisomy 10 is rare and is deleterious.

REFERENCES

1. Fryns, J.P., et al. 1991. Apparent late-onset Cockayne syndrome and interstitial deletion of the long arm of chromosome 10 (del(10)(q11.23q21.2)). *Am. J. Med. Genet.* 40: 343-344.
2. Thöny, B., et al. 1994. Chromosomal location of two human genes encoding tetrahydrobiopterin-metabolizing enzymes: 6-pyruvoyl-tetrahydropterin synthase maps to 11q22.3-q23.3, and pterin-4 α -carbinolamine dehydratase maps to 10q22. *Genomics* 19: 365-368.
3. Liu, B., et al. 2002. CARP is a novel caspase recruitment domain containing pro-apoptotic protein. *Biochem. Biophys. Res. Commun.* 293: 1396-1404.
4. Horibata, K., et al. 2004. Complete absence of Cockayne syndrome group B gene product gives rise to UV-sensitive syndrome but not Cockayne syndrome. *Proc. Natl. Acad. Sci. USA* 101: 15410-15415.
5. Teresi, R.E., et al. 2007. Cowden syndrome-affected patients with PTEN promoter mutations demonstrate abnormal protein translation. *Am. J. Hum. Genet.* 81: 756-767.
6. Cho, M.Y., et al. 2008. First report of ovarian dysgerminoma in Cowden syndrome with germline PTEN mutation and PTEN-related 10q loss of tumor heterozygosity. *Am. J. Surg. Pathol.* 32: 1258-1264.
7. Blumenthal, G.M., et al. 2008. PTEN hamartoma tumor syndromes. *Eur. J. Hum. Genet.* 16: 1289-1300.
8. Utine, G.E., et al. 2008. Kabuki syndrome and trisomy 10p. *Genet. Couns.* 19: 291-300.
9. Yin, Y., et al. 2008. PTEN: a new guardian of the genome. *Oncogene* 27: 5443-5453.

CHROMOSOMAL LOCATION

Genetic locus: FAM188A (human) mapping to 10p13; Fam188a (mouse) mapping to 2 A1.

SOURCE

FAM188A (N-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of FAM188A of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-163796 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

FAM188A (N-12) is recommended for detection of FAM188A of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other C10orf family members.

FAM188A (N-12) is also recommended for detection of FAM188A in additional species, including equine, bovine and porcine.

Suitable for use as control antibody for FAM188A siRNA (h): sc-90372, FAM188A siRNA (m): sc-108710, FAM188A shRNA Plasmid (h): sc-90372-SH, FAM188A shRNA Plasmid (m): sc-108710-SH, FAM188A shRNA (h) Lentiviral Particles: sc-90372-V and FAM188A shRNA (m) Lentiviral Particles: sc-108710-V.

Molecular Weight of FAM188A: 50 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.