

CatSper β (F-13): sc-163980

BACKGROUND

CatSper (cation channel, sperm associated proteins) are ion transport proteins located on the surface of sperm cells in the principal piece of the sperm tail. CatSper are vital to sperm motility, fertilization and cAMP-mediated calcium influx in sperm. There are four CatSper proteins in mammalian sperm, namely CatSper (or CatSper1), CatSper2, CatSper3 and CatSper4. CatSper proteins are believed to assemble into a heterotetrameric complex, forming an alkalization-activated Ca²⁺-selective channel. CatSper β is a multipass membrane protein that functions as an auxiliary protein of the CatSper channel. It contains a large extracellular domain and a number of N-glycosylation sites. CatSper β specifically interacts with CatSper1 and through this interaction plays an important role in the hyperactivated motility of sperm cells, a process that is required in the preparation of sperm for fertilization.

REFERENCES

1. Quill, T.A., et al. 2001. A voltage-gated ion channel expressed specifically in spermatozoa. *Proc. Natl. Acad. Sci. USA* 98: 12527-12531.
2. Zhang, D., et al. 2006. Association of Catsper1 or -2 with Ca(v)3.3 leads to suppression of T-type calcium channel activity. *J. Biol. Chem.* 281: 22332-22341.
3. Jin, J., et al. 2007. Catsper3 and Catsper4 are essential for sperm hyperactivated motility and male fertility in the mouse. *Biol. Reprod.* 77: 37-44.
4. Xia, J., et al. 2007. CATSPER channel-mediated Ca²⁺ entry into mouse sperm triggers a tail-to-head propagation. *Biol. Reprod.* 77: 551-559.
5. Liu, J., et al. 2007. CatSperbeta, a novel transmembrane protein in the CatSper channel complex. *J. Biol. Chem.* 282: 18945-18952.
6. Li, H.G., et al. 2007. Expression of CatSper family transcripts in the mouse testis during post-natal development and human ejaculated spermatozoa: relationship to sperm motility. *Mol. Hum. Reprod.* 13: 299-306.
7. Marquez, B., et al. 2007. Contributions of extracellular and intracellular Ca²⁺ to regulation of sperm motility: Release of intracellular stores can hyperactivate CatSper1 and CatSper2 null sperm. *Dev. Biol.* 303: 214-221.
8. Qi, H., et al. 2007. All four CatSper ion channel proteins are required for male fertility and sperm cell hyperactivated motility. *Proc. Natl. Acad. Sci. USA* 104: 1219-1223.
9. Suarez, S.S., et al. 2007. Different regulatory systems operate in the mid-piece and principal piece of the mammalian sperm flagellum. *Soc. Reprod. Fertil. Suppl.* 65: 331-334.

CHROMOSOMAL LOCATION

Genetic locus: CATSPERB (human) mapping to 14q32.12; Catsperb (mouse) mapping to 12 E.

SOURCE

CatSper β (F-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of CatSper β of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-163980 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

CatSper β (F-13) is recommended for detection of CatSper β of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

CatSper β (F-13) is also recommended for detection of CatSper β in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for CatSper β siRNA (h): sc-92375, CatSper β siRNA (m): sc-155873, CatSper β shRNA Plasmid (h): sc-92375-SH, CatSper β shRNA Plasmid (m): sc-155873-SH, CatSper β shRNA (h) Lentiviral Particles: sc-92375-V and CatSper β shRNA (m) Lentiviral Particles: sc-155873-V.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.