

E4F1 (G-15): sc-164249

BACKGROUND

E4F1 (E4F transcription factor 1), also known as E4F, is a 784 amino acid protein that localizes to both the nucleus and the cytoplasm and contains 9 C₂H₂-type zinc fingers. Expressed ubiquitously in adult and fetal tissues, E4F1 exists as a homodimer that binds DNA and is thought to act as a transcriptional repressor and may also play a role in cell survival and growth via cell cycle control. Additionally, E4F1 is thought to function as a ubiquitin ligase that mediates the ubiquitination (and subsequent degradation) of target proteins and may be involved in the p53 tumor suppressor pathway. E4F1, which may be post-translationally phosphorylated or sumoylated, is subject to proteolytic cleavage which results in the creation of a short peptide with specific DNA binding capabilities.

REFERENCES

1. Fernandes, E.R. and Rooney, R.J. 1997. The adenovirus E1A-regulated transcription factor E4F is generated from the human homolog of nuclear factor ϕ AP3. *Mol. Cell. Biol.* 17: 1890-1903.
2. Saccone, S., et al. 1998. Assignment of the E1A-regulated transcription factor E4F gene (E4F1) to human chromosome band 16p13.3 by *in situ* hybridization and somatic cell hybrids. *Cytogenet. Cell Genet.* 82: 99-100.
3. Rooney, R.J., et al. 1998. Chromosomal location and tissue expression of the gene encoding the adenovirus E1A-regulated transcription factor E4F in humans and mice. *Mamm. Genome* 9: 320-323.
4. Sandy, P., et al. 2000. p53 is involved in the p120^{E4F}-mediated growth arrest. *Oncogene* 19: 188-199.
5. Fajas, L., et al. 2000. pRB binds to and modulates the transrepressing activity of the E1A-regulated transcription factor p120^{E4F}. *Proc. Natl. Acad. Sci. USA* 97: 7738-7743.
6. Nakamura, Y., et al. 2004. E4F1, a novel estrogen-responsive gene in possible atheroprotection, revealed by microarray analysis. *Am. J. Pathol.* 165: 2019-2031.

CHROMOSOMAL LOCATION

Genetic locus: E4F1 (human) mapping to 16p13.3; E4f1 (mouse) mapping to 17 A3.3.

SOURCE

E4F1 (G-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of E4F1 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-164249 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

E4F1 (G-15) is recommended for detection of E4F1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

E4F1 (G-15) is also recommended for detection of E4F1 in additional species, including equine and bovine.

Suitable for use as control antibody for E4F1 siRNA (h): sc-93081, E4F1 siRNA (m): sc-143261, E4F1 shRNA Plasmid (h): sc-93081-SH, E4F1 shRNA Plasmid (m): sc-143261-SH, E4F1 shRNA (h) Lentiviral Particles: sc-93081-V and E4F1 shRNA (m) Lentiviral Particles: sc-143261-V.


Molecular Weight of E4F1: 83 kDa.

Positive Controls: IMR-32 nuclear extract: sc-2148.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



E4F1 (G-15): sc-164249. Western blot analysis of E4F1 expression in IMR-32 nuclear extract.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **E4F1 (D-12): sc-514718**, our highly recommended monoclonal alternative to E4F1 (G-15).