

FAM63A (S-14): sc-164337

BACKGROUND

Chromosome 1 is the largest human chromosome spanning about 260 million base pairs and making up 8% of the human genome. There are about 3,000 genes on chromosome 1, and considering the great number of genes there are also a large number of diseases associated with chromosome 1. Notably, the rare aging disease Hutchinson-Gilford progeria is associated with the LMNA gene which encodes lamin A. When defective, the LMNA gene product can build up in the nucleus and cause characteristic nuclear blebs. The mechanism of rapidly enhanced aging is unclear and is a topic of continuing exploration. The MUTYH gene is located on chromosome 1 and is partially responsible for familial adenomatous polyposis. Stickler syndrome, Parkinsons, Gaucher disease and Usher syndrome are also associated with chromosome 1. A breakpoint has been identified in 1q which disrupts the DISC1 gene and is linked to schizophrenia. Aberrations in chromosome 1 are found in a variety of cancers including head and neck cancer, malignant melanoma and multiple myeloma. The FAM63A gene product has been provisionally designated FAM63A pending further characterization.

REFERENCES

1. Watson, M.L., et al. 1990. Genomic organization of the selectin family of leukocyte adhesion molecules on human and mouse chromosome 1. *J. Exp. Med.* 172: 263-272.
2. Blackwood, D.H., et al. 2001. Schizophrenia and affective disorders— cosegregation with a translocation at chromosome 1q42 that directly disrupts brain-expressed genes: clinical and P300 findings in a family. *Am. J. Hum. Genet.* 69: 428-433.
3. Weise, A., et al. 2005. New insights into the evolution of chromosome 1. *Cytogenet. Genome Res.* 108: 217-222.
4. Gregory, S.G., et al. 2006. The DNA sequence and biological annotation of human chromosome 1. *Nature* 441: 315-321.
5. Hennah, W., et al. 2006. Genes and schizophrenia: beyond schizophrenia: the role of DISC1 in major mental illness. *Schizophr. Bull.* 32: 409-416.

CHROMOSOMAL LOCATION

Genetic locus: FAM63A (human) mapping to 1q21.3; Fam63a (mouse) mapping to 3 F2.1.

SOURCE

FAM63A (S-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of FAM63A of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-164337 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

FAM63A (S-14) is recommended for detection of FAM63A of human origin and 4930504E06Rik of mouse origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with FAM63B.

FAM63A (S-14) is also recommended for detection of FAM63A in additional species, including bovine and porcine.

Suitable for use as control antibody for FAM63A siRNA (h): sc-78698, FAM63A siRNA (m): sc-140110, FAM63A shRNA Plasmid (h): sc-78698-SH, FAM63A shRNA Plasmid (m): sc-140110-SH, FAM63A shRNA (h) Lentiviral Particles: sc-78698-V and FAM63A shRNA (m) Lentiviral Particles: sc-140110-V.

Molecular Weight of FAM63A isoform 1: 52 kDa.


Molecular Weight of FAM63A isoform 2: 36 kDa.

Molecular Weight of FAM63A isoform 3: 57 kDa.


Molecular Weight of FAM63A isoform 4: 42 kDa.

Positive Controls: FAM63A (h): 293T Lysate: sc-114955.

DATA



FAM63A (S-14): sc-164337. Western blot analysis of FAM63A expression in non-transfected: sc-117752 (A) and human FAM63A transfected: sc-114955 (B) 293T whole cell lysates.



FAM63A (S-14): sc-164337. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic and nuclear localization.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **FAM63A (F-3): sc-398287**, our highly recommended monoclonal alternative to FAM63A (S-14).