

RAP (C-18): sc-16452

BACKGROUND

Members of the LDL receptor gene family, including LDLR (low density lipoprotein receptor), LRP (low density lipoprotein related protein), Megalin (also designated GP330), VLDLR (very low density lipoprotein receptor) and ApoER2, are characterized by a cluster of cysteine-rich class A repeats, epidermal growth factor (EGF)-like repeats, YWTD repeats and an O-linked sugar domain. LRP, also designated α -2-Macroglobulin receptor, is an endocytic receptor that mediates the uptake of at least 15 ligands, including α -2-Macroglobulin and apoE. LRP is cleaved into a membrane subunit and an extracellular subunit, which remain non-covalently associated. Proper folding and trafficking of LRP is facilitated by the receptor-associated protein (RAP), a molecular chaperone. The uptake of all known ligands through LRP can be blocked by RAP, which induces a conformational change in the receptor that renders it unable to bind ligands. LRP, which is expressed in brain, liver and lung, is also implicated in Alzheimer's disease (AD), as the human LRP gene localizes to a potential AD locus on chromosome 12.

CHROMOSOMAL LOCATION

Genetic locus: LRPAP1 (human) mapping to 4p16.3; Lrpap1 (mouse) mapping to 5 B2.

SOURCE

RAP (C-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of RAP of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-16452 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

RAP (C-18) is recommended for detection of RAP of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

RAP (C-18) is also recommended for detection of RAP in additional species, including equine, canine and bovine.

Suitable for use as control antibody for RAP siRNA (h): sc-44068, RAP siRNA (m): sc-152700, RAP shRNA Plasmid (h): sc-44068-SH, RAP shRNA Plasmid (m): sc-152700-SH, RAP shRNA (h) Lentiviral Particles: sc-44068-V and RAP shRNA (m) Lentiviral Particles: sc-152700-V.


Molecular Weight of RAP: 39 kDa.

Positive Controls: A549 cell lysate: sc-2413 or RT-4 whole cell lysate: sc-364257.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



RAP (C-18): sc-16452. Western blot analysis of RAP expression in A549 (A) and RT-4 (B) whole cell lysates.



RAP (C-18): sc-16452. Immunoperoxidase staining of formalin fixed, paraffin-embedded human kidney tissue showing cytoplasmic and nuclear staining of cells in glomeruli and cells in tubules.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **RAP (E-7): sc-515625** or **RAP (7F1): sc-59675**, our highly recommended monoclonal alternatives to RAP (C-18).