

# JMJD3 (E-14): sc-164714

## BACKGROUND

JMJD3 (Jumonji domain containing 3), also known as KDM6B (lysine demethylase 6B), is a 1,679 amino acid nuclear protein that contains one JMJC domain and belongs to the highly conserved JMJC domain-containing protein family. Functioning as a histone demethylase, JMJD3 uses iron and ascorbate as cofactors to demethylate dimethylated and trimethylated Lys 27 residues of Histone H3, thereby playing an important role in the modification of the histone code. Additionally, JMJD3 regulates posterior development and is involved in the inflammatory response, specifically by mediating macrophage differentiation. JMJD3 is also thought to control the expression of neurogenesis-related proteins and, via this regulatory mechanism, may be necessary for neural commitment during early development. Two isoforms of JMJD3 exist due to alternative splicing events.

## REFERENCES

1. Cousin, P., Billotte, J., Chaubert, P. and Shaw, P. 2000. Physical map of 17p13 and the genes adjacent to p53. *Genomics* 63: 60-68.
2. Online Mendelian Inheritance in Man, OMIM<sup>™</sup>. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 611577. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
3. Xiang, Y., Zhu, Z., Han, G., Lin, H., Xu, L. and Chen, C.D. 2007. JMJD3 is a Histone H3K27 demethylase. *Cell Res.* 17: 850-857.
4. Agger, K., Cloos, P.A., Christensen, J., Pasini, D., Rose, S., Rappsilber, J., Issaeva, I., Canaani, E., Salcini, A.E. and Helin, K. 2007. UTX and JMJD3 are Histone H3K27 demethylases involved in HOX gene regulation and development. *Nature* 449: 731-734.
5. Hong, S., Cho, Y.W., Yu, L.R., Yu, H., Veenstra, T.D. and Ge, K. 2007. Identification of JMJC domain-containing UTX and JMJD3 as Histone H3 Lysine 27 demethylases. *Proc. Natl. Acad. Sci. USA* 104: 18439-18444.
6. Sen, G.L., Webster, D.E., Barragan, D.I., Chang, H.Y. and Khavari, P.A. 2008. Control of differentiation in a self-renewing mammalian tissue by the histone demethylase JMJD3. *Genes Dev.* 22: 1865-1870.
7. Li, Y., Reddy, M.A., Miao, F., Shanmugam, N., Yee, J.K., Hawkins, D., Ren, B. and Natarajan, R. 2008. Role of the Histone H3 Lysine 4 methyltransferase, SET7/9, in the regulation of NFκB dependent inflammatory genes: Relevance to diabetes and inflammation. *J. Biol. Chem.* E-published.

## CHROMOSOMAL LOCATION

Genetic locus: KDM6B (human) mapping to 17p13.1; Kdm6b (mouse) mapping to 11 B3.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## SOURCE

JMJD3 (E-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of JMJD3 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-164714 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

JMJD3 (E-14) is recommended for detection of JMJD3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other JMJD family members.

Suitable for use as control antibody for JMJD3 siRNA (h): sc-93819, JMJD3 siRNA (m): sc-146326, JMJD3 shRNA Plasmid (h): sc-93819-SH, JMJD3 shRNA Plasmid (m): sc-146326-SH, JMJD3 shRNA (h) Lentiviral Particles: sc-93819-V and JMJD3 shRNA (m) Lentiviral Particles: sc-146326-V.

Molecular Weight (predicted) of JMJD3: 180 kDa.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker<sup>™</sup> compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz<sup>™</sup> Mounting Medium: sc-24941.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.