# SANTA CRUZ BIOTECHNOLOGY, INC.

# LRRC52 (T-12): sc-164921



The Power to Question

### BACKGROUND

The leucine-rich (LRR) repeat is a 20-30 amino acid motif that forms a hydrophobic  $\alpha/\beta$  horseshoe fold, allowing it to accommodate several leucine residues within a tightly packed core. All LRR repeats contain a variable segment and a highly conserved segment, the latter of which accounts for 11 or 12 residues of the entire LRR motif. The primary function of the LRR motif is to provide a versatile structural framework to mediate the formation of protein-protein interactions. LRRs are present in a variety of proteins with diverse structure and function, including innate immunity and nervous system development. Several human diseases are associated with mutations in genes encoding LRR-containing proteins. LRRC52 (leucine-rich repeat-containing protein 52), is a 313 amino acid single-pass membrane protein that contains 5 LRR (leucine-rich) repeats, a LRRCT domain and a LRRNT domain.

# REFERENCES

- Gomi, F., et al. 2000. Molecular cloning of a novel membrane glycoprotein, pal, specifically expressed in photoreceptor cells of the retina and containing leucine-rich repeat. J. Neurosci. 20: 3206-3213.
- Kobe, B., et al. 2001. The leucine-rich repeat as a protein recognition motif. Curr. Opin. Struct. Biol. 11: 725-732.
- Hofman, P., et al. 2001. Lack of blood-brain barrier properties in microvessels of the prelaminar optic nerve head. Invest. Ophthalmol. Vis. Sci. 42: 895-901.
- 4. Hughes, J.M., et al. 2004. Vascular leucocyte adhesion molecules unaltered in the human retina in diabetes. Br. J. Ophthalmol. 88: 566-572.
- Kuiper, E.J., et al. 2004. Differential expression of connective tissue growth factor in microglia and pericytes in the human diabetic retina. Br. J. Ophthalmol. 88: 1082-1087.
- Matsushima, N., et al. 2005. Structural analysis of leucine-rich-repeat variants in proteins associated with human diseases. Cell. Mol. Life Sci. 62: 2771-2791.

## CHROMOSOMAL LOCATION

Genetic locus: LRRC52 (human) mapping to 1q24.1; Lrrc52 (mouse) mapping to 1 H2.3.

#### SOURCE

LRRC52 (T-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an N-terminal extracellular domain of LRRC52 of human origin.

#### PRODUCT

Each vial contains 200  $\mu g$  IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-164921 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

#### **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

#### APPLICATIONS

LRRC52 (T-12) is recommended for detection of LRRC52 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other LRRC family members.

Suitable for use as control antibody for LRRC52 siRNA (h): sc-78591, LRRC52 siRNA (m): sc-149092, LRRC52 shRNA Plasmid (h): sc-78591-SH, LRRC52 shRNA Plasmid (m): sc-149092-SH, LRRC52 shRNA (h) Lentiviral Particles: sc-78591-V and LRRC52 shRNA (m) Lentiviral Particles: sc-149092-V.

Molecular Weight of LRRC52: 35 kDa.

## **RECOMMENDED SECONDARY REAGENTS**

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluo-rescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

#### **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

#### PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.