Mex3a (K-16): sc-165005



The Power to Question

BACKGROUND

Mex-3 (muscle excess protein-3) is a translational regulator in *Caenorhabditis elegans* that participates in maintaining the germline totipotency and in specification of posterior blastomeres in early embryos. In humans, four evolutionarily conserved Mex-3 homologs exist, namely Mex3a, Mex3b, Mex3c and Mex3d. These proteins comprise a family of RNA binding phosphoproteins which each contain two tandemly repeated KH (nuclear ribonucleoprotein K homology) domains and one C-terminal RING finger motif. In addition, the Mex-3 homolog family of proteins shuttle between the nucleus and the cytoplasm through the CRM1-dependent export pathway and may play a role regulating post-transcriptional events. Mex3a is a 520 amino acid protein that localizes to both the nucleus and the cytoplasm and contains one RING-type zinc finger and two KH domains through which it functions as an RNA-binding protein.

REFERENCES

- Fillman, C. and Lykke-Andersen, J. 2005. RNA decapping inside and outside of processing bodies. Curr. Opin. Cell Biol. 17: 326-331.
- Buchet-Poyau, K., Courchet, J., Le Hir, H., Séraphin, B., Scoazec, J.Y., Duret, L., Domon-Dell, C., Freund, J.N. and Billaud, M. 2007. Identification and characterization of human Mex-3 proteins, a novel family of evolutionarily conserved RNA-binding proteins differentially localized to processing bodies. Nucleic Acids Res. 35: 1289-1300.
- 3. Courchet, J., Buchet-Poyau, K., Potemski, A., Brès, A., Jariel-Encontre, I. and Billaud, M. 2008. Interaction with 14-3-3 adaptors regulates the sorting of hMex3b RNA-binding protein to distinct classes of RNA granules. J. Biol. Chem. 283: 32131-32142.
- 4. Online Mendelian Inheritance in Man, OMIM™. 2009. Johns Hopkins University, Baltimore, MD. MIM Number: 611007. World Wide Web URL: http://www.ncbi.nlm.nih.gov/omim/

CHROMOSOMAL LOCATION

Genetic locus: MEX3A (human) mapping to 1q22; Mex3a (mouse) mapping to 3 F1.

SOURCE

Mex3a (K-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Mex3a of human origin.

PRODUCT

Each vial contains 200 μg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-165005 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-165005 X, 200 $\mu g/0.1$ ml.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Mex3a (K-16) is recommended for detection of Mex3a of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other Mex3 family members.

Suitable for use as control antibody for Mex3a siRNA (h): sc-88413, Mex3a siRNA (m): sc-149395, Mex3a shRNA Plasmid (h): sc-88413-SH, Mex3a shRNA Plasmid (m): sc-149395-SH, Mex3a shRNA (h) Lentiviral Particles: sc-88413-V and Mex3a shRNA (m) Lentiviral Particles: sc-149395-V.

Mex3a (K-16) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of Mex3a: 54 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3800 fax 831.457.3801 **Europe** +00800 4573 8000 49 6221 4503 0 **www.scbt.com**