

Pur α (T-13): sc-165318

BACKGROUND

The Pur protein family consists of four members: Pur α , Pur β and two isoforms of Pur γ . Pur α , a protein strongly conserved throughout evolution, is a single-stranded (ss) DNA- and RNA-binding protein that contains three conserved signature repeats, which have an affinity for guanosine-rich motifs, specifically the (GGN)_n sequence, PUR element. The ubiquitously expressed Pur α protein is involved in many cellular processes, including transcriptional regulation, the cell cycle, oncogenic transformation and postnatal brain development. Pur α binds HIV-1 TAR RNA and activates HIV-1 transcription. Pur α also appears to play a role in the progression of Alzheimer's disease, prostate cancer, fragile X-associated tremor/ataxia syndrome and JC virus. Targeting of Pur α may potentially lead to promising therapeutic approaches for various diseases.

REFERENCES

1. Haas, S., Thatikunta, P., Steplewski, A., Johnson, E.M., Khalili, K. and Amini, S. 1995. A 39-kD DNA-binding protein from mouse brain stimulates transcription of myelin basic protein gene in oligodendrocytic cells. *J. Cell Biol.* 130: 1171-1179.
2. Chepenik, L.G., Tretiakova, A.P., Krachmarov, C.P., Johnson, E.M. and Khalili, K. 1998. The single-stranded DNA binding protein, Pur α , binds HIV-1 TAR RNA and activates HIV-1 transcription. *Gene* 210: 37-44.
3. Johnson, E.M. 2003. The Pur protein family: clues to function from recent studies on cancer and AIDS. *Anticancer Res.* 23: 2093-2100.
4. Zeng, L.H., Fujimoto, T., Kumamaru, E., Irie, Y., Miki, N. and Kuo, C.H. 2004. Characterization of novel Pur α -binding proteins in mouse brain. *Neurochem. Int.* 45: 753-758.
5. Wortman, M.J., Johnson, E.M. and Bergemann, A.D. 2005. Mechanism of DNA binding and localized strand separation by Pur α and comparison with Pur family member, Pur β . *Biochim. Biophys. Acta* 1743: 64-78.
6. Darbinian, N., White, M.K. and Khalili, K. 2006. Regulation of the Pur α promoter by E2F-1. *J. Cell. Biochem.* 99: 1052-1063.
7. Jin, P., Duan, R., Qurashi, A., Qin, Y., Tian, D., Rosser, T.C., Liu, H., Feng, Y. and Warren, S.T. 2007. Pur α binds to rCGG repeats and modulates repeat-mediated neurodegeneration in a *Drosophila* model of fragile X tremor/ataxia syndrome. *Neuron* 55: 556-564.
8. Wang, L.G., Johnson, E.M., Kinoshita, Y., Babb, J.S., Buckley, M.T., Liebes, L.F., Melamed, J., Liu, X.M., Kurek, R., Ossowski, L. and Ferrari, A.C. 2008. Androgen receptor overexpression in prostate cancer linked to Pur α loss from a novel repressor complex. *Cancer Res.* 68: 2678-2688.

CHROMOSOMAL LOCATION

Genetic locus: PURA (human) mapping to 5q31.2; Pura (mouse) mapping to 18 B2.

SOURCE

Pur α (T-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Pur α of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-165318 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-165318 X, 200 μ g/0.1 ml.

APPLICATIONS

Pur α (T-13) is recommended for detection of Pur α of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Pur α (T-13) is also recommended for detection of Pur α in additional species, including canine, bovine, porcine and avian.

Suitable for use as control antibody for Pur α siRNA (h): sc-91623, Pur α siRNA (m): sc-152591, Pur α shRNA Plasmid (h): sc-91623-SH, Pur α shRNA Plasmid (m): sc-152591-SH, Pur α shRNA (h) Lentiviral Particles: sc-91623-V and Pur α shRNA (m) Lentiviral Particles: sc-152591-V.

Pur α (T-13) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of Pur α : 39 kDa.

Positive Controls: human skeletal muscle extract: sc-363776.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



Try **Pur α (80-L): sc-130397**, our highly recommended monoclonal alternative to Pur α (T-13).