

# Rab11-FIP4 (S-14): sc-165331

## BACKGROUND

The Ras superfamily of GTPases can be subdivided into the Ras, Rho/Rac, Sar, Rab, Arf, Rap and Ran subfamilies, all of which control multiple aspects of cell function, including cytoskeletal rearrangement, nuclear signaling and cell growth. Members of the Ras protein superfamily are regulated by a variety of GTPase-interaction proteins that control GTPase function. Rab11-FIP4 (RAB11 family interacting protein 4 (class II)), also known as arfophilin-2, is a 637 amino acid peripheral membrane protein that localizes to the midbody and cleavage furrow during cytokinesis. Highly expressed in testis with weak expression found in other tissues, Rab11-FIP4 plays a role in endocytic trafficking and cytokinesis, and exists as a homodimer that forms a complex with ARF6 and Rab11. Rab11-FIP4 contains one EF-hand domain, an RBD-FIP domain and exists as two alternatively spliced isoforms.

## REFERENCES

1. Nagase, T., Nakayama, M., Nakajima, D., Kikuno, R. and Ohara, O. 2001. Prediction of the coding sequences of unidentified human genes. XX. The complete sequences of 100 new cDNA clones from brain which code for large proteins *in vitro*. DNA Res. 8: 85-95.
2. Hales, C.M., Griner, R., Hobby-Henderson, K.C., Dorn, M.C., Hardy, D., Kumar, R., Navarre, J., Chan, E.K., Lapierre, L.A. and Goldenring, J.R. 2001. Identification and characterization of a family of Rab11-interacting proteins. J. Biol. Chem. 276: 39067-39075.
3. Wallace, D.M., Lindsay, A.J., Hendrick, A.G. and McCaffrey, M.W. 2002. Rab11-FIP4 interacts with Rab11 in a GTP-dependent manner and its over-expression condenses the Rab11 positive compartment in HeLa cells. Biochem. Biophys. Res. Commun. 299: 770-779.
4. Fielding, A.B., Schonteich, E., Matheson, J., Wilson, G., Yu, X., Hickson, G.R., Srivastava, S., Baldwin, S.A., Prekeris, R. and Gould, G.W. 2005. Rab11-FIP3 and FIP4 interact with Arf6 and the exocyst to control membrane traffic in cytokinesis. EMBO J. 24: 3389-3399.
5. Online Mendelian Inheritance in Man, OMIM<sup>™</sup>. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 179508; World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

## CHROMOSOMAL LOCATION

Genetic locus: RAB11FIP4 (human) mapping to 17q11.2; Rab11fip4 (mouse) mapping to 11 B5.

## SOURCE

Rab11-FIP4 (S-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Rab11-FIP4 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-165331 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

Rab11-FIP4 (S-14) is recommended for detection of Rab11-FIP4 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other Rab11-FIP family members.

Suitable for use as control antibody for Rab11-FIP4 siRNA (h): sc-93812, Rab11-FIP4 siRNA (m): sc-152663, Rab11-FIP4 shRNA Plasmid (h): sc-93812-SH, Rab11-FIP4 shRNA Plasmid (m): sc-152663-SH, Rab11-FIP4 shRNA (h) Lentiviral Particles: sc-93812-V and Rab11-FIP4 shRNA (m) Lentiviral Particles: sc-152663-V.

Molecular Weight of Rab11-FIP4: 72 kDa.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker<sup>™</sup> compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz<sup>™</sup> Mounting Medium: sc-24941.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.