

SPNS2 (N-14): sc-165574

BACKGROUND

SPNS2 (spinster homolog 2) is a 549 amino acid multi-pass membrane protein that belongs to the spinster subfamily of major facilitator proteins. The gene encoding SPNS2 maps to human chromosome 17, which comprises over 2.5% of the human genome and encodes over 1,200 genes. Two key tumor suppressor genes are associated with chromosome 17, namely, p53 and BRCA1. Tumor suppressor p53 is necessary for maintenance of cellular genetic integrity by moderating cell fate through DNA repair versus cell death. Malfunction or loss of p53 expression is associated with malignant cell growth and Li-Fraumeni syndrome. Like p53, BRCA1 is directly involved in DNA repair, though specifically it is recognized as a genetic determinant of early onset breast cancer and predisposition to cancers of the ovary, colon, prostate gland and fallopian tubes.

REFERENCES

- Zhang, Z.Y., et al. 2006. Expression of MAC30 in rectal cancers with or without preoperative radiotherapy. *Oncology* 71: 259-265.
- Wilcox, C.B., et al. 2007. Coordinate up-regulation of TMEM97 and cholesterol biosynthesis genes in normal ovarian surface epithelial cells treated with progesterone: implications for pathogenesis of ovarian cancer. *BMC Cancer* 7: 223.
- Nusbaum, R., et al. 2006-2007. Susceptibility to breast cancer: hereditary syndromes and low penetrance genes. *Breast Dis.* 27: 21-50.
- Ropolo, A., et al. 2007. The pancreatitis-induced vacuole membrane protein 1 triggers autophagy in mammalian cells. *J. Biol. Chem.* 282: 37124-37133.
- Tai, Y.C., et al. 2007. Breast cancer risk among male BRCA1 and BRCA2 mutation carriers. *J. Natl. Cancer Inst.* 99: 1811-1814.
- Yan, J., et al. 2007. BLIMP1 regulates cell growth through repression of p53 transcription. *Proc. Natl. Acad. Sci. USA* 104: 1841-1846.
- Moparthy, S.B., et al. 2007. Expression of MAC30 protein is related to survival and biological variables in primary and metastatic colorectal cancers. *Int. J. Oncol.* 30: 91-95.
- Saueremann, M., et al. 2008. Reduced expression of vacuole membrane protein 1 affects the invasion capacity of tumor cells. *Oncogene* 27: 1320-1326.
- Kawahara, A., et al. 2009. The sphingolipid transporter spns2 functions in migration of zebrafish myocardial precursors. *Science* 323: 524-527.

CHROMOSOMAL LOCATION

Genetic locus: SPNS2 (human) mapping to 17p13.3; Spns2 (mouse) mapping to 11 B4.

SOURCE

SPNS2 (N-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of SPNS2 of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-165574 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

SPNS2 (N-14) is recommended for detection of SPNS2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with SPNS1 or SPNS3.

SPNS2 (N-14) is also recommended for detection of SPNS2 in additional species, including bovine and porcine.

Suitable for use as control antibody for SPNS2 siRNA (h): sc-106749, SPNS2 shRNA Plasmid (h): sc-106749-SH and SPNS2 shRNA (h) Lentiviral Particles: sc-106749-V.

Molecular Weight of SPNS2: 58 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.