

SURF-2 (T-13): sc-165627

BACKGROUND

SURF-2 (surfeit locus protein 2) is a 256 amino acid protein that belongs to the SURF2 family and interacts with β -1,4-Gal-T3, uPAR and WDR20. The gene that encodes SURF-2 is located in the surfeit gene cluster, which is a group of very tightly linked genes that do not share sequence similarity. The SURF-2 gene maps to human chromosome 9q34.2 and shares a bidirectional promoter with SURF1, which is located on the opposite strand. The intergenic region between the SURF-1 and SURF-2 genes is expected to have bidirectional promoter activity, as is found in mouse. Although this region lacks a TATA box, it is GC-rich. Housing over 900 genes, human chromosome 9 comprises nearly 4% of the human genome. Hereditary hemorrhagic telangiectasia, which is characterized by harmful vascular defects, and familial dysautonomia, are both associated with chromosome 9.

REFERENCES

1. Online Mendelian Inheritance in Man, OMIM[™]. 1989. Johns Hopkins University, Baltimore, MD. MIM Number: 185630. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
2. Yon, J., et al. 1993. The organization and conservation of the human Surfeit gene cluster and its localization telomeric to the c-abl and can proto-oncogenes at chromosome band 9q34.1. *Hum. Mol. Genet.* 2: 237-240.
3. Lennard, A., et al. 1994. The Surf-1 and Surf-2 genes and their essential bidirectional promoter elements are conserved between mouse and human. *DNA Cell Biol.* 13: 1117-1126.
4. Duhig, T., et al. 1998. The human Surfeit locus. *Genomics* 52: 72-78.
5. Cottin, V., et al. 2007. Pulmonary vascular manifestations of hereditary hemorrhagic telangiectasia (rendu-osler disease). *Respiration* 74: 361-378.
6. Zeitz, M.J., et al. 2009. Organization of the amplified type I interferon gene cluster and associated chromosome regions in the interphase nucleus of human osteosarcoma cells. *Chromosome Res.* 17: 305-319.
7. Gold-von Simson, G., et al. 2009. Kinetin in familial dysautonomia carriers: implications for a new therapeutic strategy targeting mRNA splicing. *Pediatr. Res.* 65: 341-346.
8. Axelrod, F.B., et al. 2010. Neuroimaging supports central pathology in familial dysautonomia. *J. Neurol.* 257: 198-206.

CHROMOSOMAL LOCATION

Genetic locus: SURF2 (human) mapping to 9q34.2; Surf2 (mouse) mapping to 2 A3.

SOURCE

SURF-2 (T-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of SURF-2 of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-165627 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

SURF-2 (T-13) is recommended for detection of SURF-2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other SURF family members.

Suitable for use as control antibody for SURF-2 siRNA (h): sc-92780, SURF-2 siRNA (m): sc-153934, SURF-2 shRNA Plasmid (h): sc-92780-SH, SURF-2 shRNA Plasmid (m): sc-153934-SH, SURF-2 shRNA (h) Lentiviral Particles: sc-92780-V and SURF-2 shRNA (m) Lentiviral Particles: sc-153934-V.

Molecular Weight of SURF-2: 30 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.