

# UNC45A (T-12): sc-165863

UNC45A (UNC-45 homolog A), also known as SMAP1 (smooth muscle cell-associated protein 1) or GCUNC45, is a 944 amino acid protein that localizes to the perinuclear region of the cytoplasm and contains three TPR repeats. Expressed in spinal cord, bone marrow, peripheral blood leukocytes, trachea, stomach and adrenal gland, UNCA45A functions as a chaperone for HSP 90, effectively facilitating the association of HSP 90 with PR (progesterone receptor). Additionally, UNC45A is thought to be necessary for proper folding of myosin and is required for normal myotube formation during muscle cell development, thereby playing an important role in muscle formation. Overexpression of UNC45A is associated with rapid cell proliferation and motility, especially that observed in ovarian cancer, suggesting a role for UNC45A in tumorigenesis. Multiple isoforms of UNC45A exist due to alternative splicing events.

## REFERENCES

1. Price, M.G., Landsverk, M.L., Barral, J.M. and Epstein, H.F. 2002. Two mammalian UNC-45 isoforms are related to distinct cytoskeletal and muscle-specific functions. *J. Cell Sci.* 115: 4013-4023.
2. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 611219. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
3. Chadli, A., Graham, J.D., Abel, M.G., Jackson, T.A., Gordon, D.F., Wood, W.M., Felts, S.J., Horwitz, K.B. and Toft, D. 2006. GCUNC45 is a novel regulator for the progesterone receptor/HSP 90 chaperoning pathway. *Mol. Cell. Biol.* 26: 1722-1730.
4. Bazzaro, M., Santillan, A., Lin, Z., Tang, T., Lee, M.K., Bristow, R.E., Shih, I.e.M. and Roden, R.B. 2007. Myosin II cochaperone general cell UNC45 overexpression is associated with ovarian cancer, rapid proliferation, and motility. *Am. J. Pathol.* 171: 1640-1649.
5. Chadli, A., Bruinsma, E.S., Stensgard, B. and Toft, D. 2008. Analysis of HSP 90 cochaperone interactions reveals a novel mechanism for TPR protein recognition. *Biochemistry* 47: 2850-2857.
6. Anderson, M.J., Pham, V.N., Vogel, A.M., Weinstein, B.M. and Roman, B.L. 2008. Loss of UNC45A precipitates arteriovenous shunting in the aortic arches. *Dev. Biol.* 318: 258-267.
7. Chadli, A., Felts, S.J. and Toft, D.O. 2008. GCUNC45 is the first HSP 90 cochaperone to show  $\alpha/\beta$  isoform specificity. *J. Biol. Chem.* 283: 9509-9512.
8. Liu, L., Sriakulam, R. and Winkelmann, D.A. 2008. UNC45 activates HSP 90-dependent folding of the myosin motor domain. *J. Biol. Chem.* 283: 13185-13193.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## CHROMOSOMAL LOCATION

Genetic locus: UNC45A (human) mapping to 15q26.1; Unc45a (mouse) mapping to 7 D3.

## SOURCE

UNC45A (T-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of UNC45A of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-165863 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

UNC45A (T-12) is recommended for detection of UNC45A of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with UNC45B.

Suitable for use as control antibody for UNC45A siRNA (h): sc-90291, UNC45A siRNA (m): sc-154919, UNC45A shRNA Plasmid (h): sc-90291-SH, UNC45A shRNA Plasmid (m): sc-154919-SH, UNC45A shRNA (h) Lentiviral Particles: sc-90291-V and UNC45A shRNA (m) Lentiviral Particles: sc-154919-V.

Molecular Weight of UNC45A: 103 kDa.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.