

ZFYVE19 (T-16): sc-165941

BACKGROUND

Zinc finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. ZFYVE19 (zinc finger, FYVE domain containing 19), also known as MPFYVE (MLL partner containing FYVE domain), is a 471 amino acid protein that contains one FYVE-type zinc finger. Expressed in heart, brain, kidney, skeletal muscle and liver, ZFYVE19 may participate in transcriptional regulation events within the cell. Defects in the gene encoding ZFYVE19 are associated with acute myeloblastic leukemia (AML), a rapidly progressing cancer of the myeloid line of white blood cells that is characterized by fever, anemia, bone pain, shortness of breath and frequent infections. Three isoforms of ZFYVE19 exist due to alternative splicing events.

REFERENCES

- Hayette, S., Tigaud, I., Vanier, A., Martel, S., Corbo, L., Charrin, C., Beillard, E., Deleage, G., Magaud, J.P. and Rimokh, R. 2000. AF15q14, a novel partner gene fused to the MLL gene in an acute myeloid leukaemia with a t(11;15)(q23;q14). *Oncogene* 19: 4446-4450.
- Shibuya, N., Taki, T., Mugishima, H., Chin, M., Tsuchida, M., Sako, M., Kawa, K., Ishii, E., Miura, I., Yanagisawa, M. and Hayashi, Y. 2001. t(10;11)-acute leukemias with MLL-AF10 and MLL-ABI1 chimeric transcripts: specific expression patterns of ABI1 gene in leukemia and solid tumor cell lines. *Genes Chromosomes Cancer* 32: 1-10.
- Ono, R., Taki, T., Taketani, T., Taniwaki, M., Kobayashi, H. and Hayashi, Y. 2002. LCX, leukemia-associated protein with a CXXC domain, is fused to MLL in acute myeloid leukemia with trilineage dysplasia having t(10;11)(q22;q23). *Cancer Res.* 62: 4075-4080.
- Chinwalla, V., Chien, A., Odero, M., Neilly, M.B., Zeleznik-Le, N.J. and Rowley, J.D. 2003. A t(11;15) fuses MLL to two different genes, AF15q14 and a novel gene MPFYVE on chromosome 15. *Oncogene* 22: 1400-1410.
- Kuefer, M.U., Chinwalla, V., Zeleznik-Le, N.J., Behm, F.G., Naeve, C.W., Rakestraw, K.M., Mukatira, S.T., Raimondi, S.C. and Morris, S.W. 2003. Characterization of the MLL partner gene AF15q14 involved in t(11;15)(q23;q14). *Oncogene* 22: 1418-1424.
- Olsen, J.V., Blagoev, B., Gnad, F., Macek, B., Kumar, C., Mortensen, P. and Mann, M. 2006. Global, *in vivo*, and site-specific phosphorylation dynamics in signaling networks. *Cell* 127: 635-648.

CHROMOSOMAL LOCATION

Genetic locus: ZFYVE19 (human) mapping to 15q15.1; Zfyve19 (mouse) mapping to 2 E5.

SOURCE

ZFYVE19 (T-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of ZFYVE19 of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-165941 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

ZFYVE19 (T-16) is recommended for detection of ZFYVE19 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other ZFYVE family members.

Suitable for use as control antibody for ZFYVE19 siRNA (h): sc-90097, ZFYVE19 siRNA (m): sc-155604, ZFYVE19 shRNA Plasmid (h): sc-90097-SH, ZFYVE19 shRNA Plasmid (m): sc-155604-SH, ZFYVE19 shRNA (h) Lentiviral Particles: sc-90097-V and ZFYVE19 shRNA (m) Lentiviral Particles: sc-155604-V.

Molecular Weight of ZFYVE19: 48 kDa.

Positive Controls: PLC/PRF/5 whole cell lysate.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.