ZNF23 (E-15): sc-165950



The Power to Question

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZNF23 (zinc finger protein 23), also known as KOX16, ZNF359, ZNF612 or Zfp612, is a 643 amino acid protein that localizes to the nucleus and contains 17 $\rm C_2H_2$ -type zinc fingers and a KRAB domain. One of several members of the Krüppel $\rm C_2H_2$ -type zinc-finger protein family, ZNF23 is thought to be involved in transcriptional regulation events. The gene encoding ZNF23 maps to human chromosome 16, which encodes over 900 genes and comprises nearly 3% of the human genome.

REFERENCES

- Cannizzaro, L.A., et al. 1993. Human zinc finger gene ZNF23 (Kox16) maps to a zinc finger gene cluster on chromosome 16q22, and ZNF32 (Kox30) to chromosome region 10q23-q24. Hum. Genet. 91: 383-385.
- Rousseau-Merck, M.F., et al. 1994. Chromosomal localization of 9 KOX zinc finger genes: physical linkages suggest clustering of KOX genes on chromosomes 12, 16, and 19. Hum. Genet. 92: 583-587.
- 3. Sun, Y., et al. 2003. The KRAB domain of zinc finger gene ZNF268: a potential transcriptional repressor. IUBMB Life 55: 127-131.
- 4. Rousseau-Merck, et al. 2003. The KOX zinc finger genes: genome wide mapping of 368 ZNF PAC clones with zinc finger gene clusters predominantly in 23 chromosomal loci are confirmed by human sequences annotated in EnsEMBL. Cytogenet. Genome Res. 98: 147-153.
- Nakamura, M., et al. 2004. A novel subfamily of zinc finger genes involved in embryonic development. J. Cell. Biochem. 93: 887-895.
- Englbrecht, C.C., et al. 2004. Conservation, diversification and expansion of C₂H₂ zinc finger proteins in the *Arabidopsis thaliana* genome. BMC Genomics 5: 39.
- 7. Li, Y., et al. 2006. A novel zinc-finger protein ZNF436 suppresses transcriptional activities of AP-1 and SRE. Mol. Biol. Rep. 33: 287-294.
- 8. Zhong, Z., et al. 2007. Identification of a novel human zinc finger gene, ZNF438, with transcription inhibition activity. J. Biochem. Mol. Biol. 40: 517-524.
- 9. O'Geen, H., et al. 2007. Genome-wide analysis of KAP1 binding suggests autoregulation of KRAB-ZNFs. PLoS Genet. 3: e89.

CHROMOSOMAL LOCATION

Genetic locus: ZNF23 (human) mapping to 16q22.2.

SOURCE

ZNF23 (E-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of ZNF23 of human origin.

PRODUCT

Each vial contains 200 μg IgG in 1.0 ml of PBS with <0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-165950 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

ZNF23 (E-15) is recommended for detection of ZNF23 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other ZNF family members.

Suitable for use as control antibody for ZNF23 siRNA (h): sc-93526, ZNF23 shRNA Plasmid (h): sc-93526-SH and ZNF23 shRNA (h) Lentiviral Particles: sc-93526-V.

Molecular Weight of ZNF23: 73 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 Fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com