

G_β (C-8): sc-166064

BACKGROUND

Heterotrimeric G proteins function to relay information from cell surface receptors to intracellular effectors. Each of a very broad range of receptors specifically detects an extracellular stimulus (i.e., a photon, pheromone, odorant, hormone or neurotransmitter), while the effectors (e.g., adenylyl cyclase), which act to generate one or more intracellular messengers, are less numerous. Each subunit of the G protein complex is encoded by a member of one of three corresponding gene families (α , β , γ). In mammals, there are five different members of the β -subunit family. The β subunits of the G proteins are important regulators of G protein α subunits as well as of certain signal transduction receptors and effectors. In contrast to G _{β 1-4}, which are at least 83% homologous, G _{β 5} is only 50% homologous to the other β subunits. Human G _{β 5} is expressed at high levels in brain, pancreas, kidney, and heart.

REFERENCES

- Gautam, N., et al. 1990. G protein diversity is increased by associations with a variety of γ subunits. *Proc. Natl. Acad. Sci. USA* 87: 7973-7977.
- Simon, M.I., et al. 1991. Diversity of G proteins in signal transduction. *Science* 252: 802-808.
- von Weizsacker, E., et al. 1992. Diversity among the β subunits of heterotrimeric GTP-binding proteins: characterization of a novel β subunit cDNA. *Biochem. Biophys. Res. Commun.* 183: 350-356.
- Kleuss, C., et al. 1992. Different β subunits determine G protein interaction with transmembrane receptors. *Nature* 358: 424-426.
- Blank, J.L., et al. 1992. Activation of cytosolic phosphoinositide phospholipase C by G protein $\beta\gamma$ subunits. *J. Biol. Chem.* 267: 23069-23075.
- Íñiguez-Lluhi, J.A., et al. 1992. G protein $\beta\gamma$ subunits synthesized in Sf9 cells. *J. Biol. Chem.* 267: 23409-23417.
- Cali, J.J., et al. 1992. Selective tissue distribution of G protein γ subunits, including a new form of the γ subunits identified by cDNA cloning. *J. Biol. Chem.* 267: 24023-24027.
- Conklin, B.R., et al. 1993. Structural elements of G α subunits that interact with G $\beta\gamma$ receptors, and effectors. *Cell* 73: 631-641.
- Kleuss, C., et al. 1993. Selectivity in signal transduction determined by γ subunits of heterotrimeric G proteins. *Science* 259: 832-834.

SOURCE

G _{β} (C-8) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 302-340 at the C-terminus of G _{β} of mouse origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-166064 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

G _{β} (C-8) is recommended for detection of G _{β 1}, G _{β 2}, G _{β 3} and G _{β 4} of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Molecular Weight of G _{β} : 36 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204, AMJ2-C8 whole cell lysate: sc-364366 or NIH/3T3 whole cell lysate: sc-2210.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgG κ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



G _{β} (C-8): sc-166064. Western blot analysis of G _{β} expression in Jurkat (A), U-251-MG (B), NIH/3T3 (C), AMJ2-C8 (D), C6 (E) and NRK (F) whole cell lysates.



G _{β} (C-8): sc-166064. Immunoperoxidase staining of formalin fixed, paraffin-embedded human gall bladder tissue showing cytoplasmic staining of glandular cells.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.