

V-ATPase B2 (C-9): sc-166122

BACKGROUND

Vacuolar-type H⁺-ATPase (V-ATPase) is a multisubunit enzyme responsible for acidification of eukaryotic intracellular organelles. V-ATPases pump protons against an electrochemical gradient, while F-ATPases reverse the process, thereby synthesizing ATP. A peripheral V₁ domain, which is responsible for ATP hydrolysis and an integral V₀ domain, which is responsible for proton translocation, compose V-ATPase. Nine subunits (A-H) make up the V₁ domain and five subunits (a, d, c, c' and c'') make up the V₀ domain. Like F-ATPase, V-ATPase most likely operates through a rotary mechanism. The V-ATPase V₁ B subunit exists as two isoforms. In the inner ear, the V-ATPase B1 isoform functions in proton secretion and is required to maintain proper endolymph pH and normal auditory function. The gene encoding the human V-ATPase B1 isoform maps to chromosome 2p13.3. Mutations in this gene cause distal renal tubular acidosis associated with sensorineural deafness. The V-ATPase B2 isoform is expressed in kidney and is the only B isoform expressed in osteoclasts. The gene encoding the human V-ATPase B2 isoform maps to chromosome 8p21.3.

CHROMOSOMAL LOCATION

Genetic locus: ATP6V1B2 (human) mapping to 8p21.3; Atp6v1b2 (mouse) mapping to 8 B3.3.

SOURCE

V-ATPase B2 (C-9) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 2-38 at the N-terminus of V-ATPase B2 of human origin.

PRODUCT

Each vial contains 200 µg IgG₃ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-166122 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

V-ATPase B2 (C-9) is recommended for detection of V-ATPase B2 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

V-ATPase B2 (C-9) is also recommended for detection of V-ATPase B2 in additional species, including canine and bovine.


Suitable for use as control antibody for V-ATPase B2 siRNA (h): sc-43517, V-ATPase B2 siRNA (m): sc-43518, V-ATPase B2 shRNA Plasmid (h): sc-43517-SH, V-ATPase B2 shRNA Plasmid (m): sc-43518-SH, V-ATPase B2 shRNA (h) Lentiviral Particles: sc-43517-V and V-ATPase B2 shRNA (m) Lentiviral Particles: sc-43518-V.

Molecular Weight of V-ATPase B2: 56-58 kDa.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



V-ATPase B2 (C-9): sc-166122. Western blot analysis of V-ATPase B2 expression in SK-N-SH (A), HeLa (B), c4 (C), RAW 264.7 (D), L8 (E) and C6 (F) whole cell lysates.



V-ATPase B2 (C-9): sc-166122. Immunofluorescence staining of formalin-fixed Hep G2 cells showing cytoplasmic and cytoplasmic vesicles localization.

SELECT PRODUCT CITATIONS

- Yang, N.D., et al. 2014. Artesunate induces cell death in human cancer cells via enhancing lysosomal function and lysosomal degradation of ferritin. *J. Biol. Chem.* 289: 33425-33441.
- Shi, Y., et al. 2015. Critical role of CAV1/caveolin-1 in cell stress responses in human breast cancer cells via modulation of lysosomal function and autophagy. *Autophagy* 11: 769-784.
- Hirata, H., et al. 2021. PMEPA1 and NEDD4 control the proton production of osteoclasts by regulating vesicular trafficking. *FASEB J.* 35: e21281.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.