SANTA CRUZ BIOTECHNOLOGY, INC.

SR-A (B-9): sc-166139



BACKGROUND

The macrophage class A scavenger receptor (SR-A) mediates the uptake of modified low density lipoprotein (LDL). The gene encoding human SR-A maps to chromosome 8 and gives rise to two alternatively spliced isoforms, type I and II (SR-AI and SR-AII), which were originally cloned from the phorbol ester-treated human monocytic cell line THP-1. Both isoforms contain six domains: cytoplasmic (I), membrane-spanning (II), spacer (III), α -helical coiled-coil (IV), collagen-like (V) and a type-specific C-terminal (VI). Domain IV is essential for the trimerization of SR-A, whereas domain V is essential for the wide range of ligand recognition. SR-A is expressed in liver, placenta and brain. Both SR-AI and SR-AII mediate the uptake of LDLs in atherosclerotic lesions. A third isoform, SR-AIII, is unable to uptake LDLs and acts as a dominant negative isoform to possibly protect cells found in advanced atherosclerotic lesions. SR-A plays a role not only in many macrophage-associated pathological pro-cesses, including atherosclerosis and Alzheimer's disease, but also in host defense and as an adhesion molecule.

CHROMOSOMAL LOCATION

Genetic locus: MSR1 (human) mapping to 8p22; Msr1 (mouse) mapping to 8 A4.

SOURCE

SR-A (B-9) is a mouse monoclonal antibody raised against amino acids 61-250 of SR-A of human origin.

PRODUCT

Each vial contains 200 μ g lgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

SR-A (B-9) is available conjugated to agarose (sc-166139 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-166139 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-166139 PE), fluorescein (sc-166139 FITC), Alexa Fluor[®] 488 (sc-166139 AF488), Alexa Fluor[®] 546 (sc-166139 AF546), Alexa Fluor[®] 594 (sc-166139 AF594) or Alexa Fluor[®] 647 (sc-166139 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-166139 AF680) or Alexa Fluor[®] 790 (sc-166139 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

APPLICATIONS

SR-A (B-9) is recommended for detection of SR-A of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for SR-A siRNA (h): sc-44116, SR-A siRNA (m): sc-40188, SR-A shRNA Plasmid (h): sc-44116-SH, SR-A shRNA Plasmid (m): sc-40188-SH, SR-A shRNA (h) Lentiviral Particles: sc-44116-V and SR-A shRNA (m) Lentiviral Particles: sc-40188-V.

Molecular Weight of glycosylated SR-A: 75 kDa.

Molecular Weight of SR-A isoforms: 50/40/43 kDa.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA





SR-A (B-9): sc-166139. Western blot analysis of SR-A expression in untreated (**A**) and chemically-treated (**B**, **C**, **D**) HEK293T whole cell lysates. β -Actin (C4): sc-47778 used as loading control. Detection reagent used: m-IgG Fc BP-HRP: sc-525409 SR-A (B-9): sc-166139. Western blot analysis of SR-A expression in non-transfected: sc-117752 (A) and human SR-A transfected: sc-111501 (B) 293T whole cell lysates.

SELECT PRODUCT CITATIONS

- Ren, Z., et al. 2020. Anti-glycolipid disorder effect of epigallocatechin-3gallate on high-fat diet and STZ-induced T2DM in mice. Mol. Med. Rep. 21: 2475-2483.
- Zhu, X., et al. 2021. SR-A1 prevents obesity-associated blood pressure elevation through suppressing overproduction of VEGF-B in macrophages. Cardiovasc. Res. 117: 547-560.
- 3. Cheng, W.L., et al. 2021. ALK7 acts as a positive regulator of macrophage activation through down-regulation of PPAR γ expression. J. Atheroscler. Thromb. 28: 375-384.
- Cheng, W.L., et al. 2021. PAK1 silencing attenuated proinflammatory macrophage activation and foam cell formation by increasing PPARγ expression. Oxid. Med. Cell. Longev. 2021: 6957900.
- Turati, J., et al. 2022. A metabotropic glutamate receptor 3 (mGlu3R) isoform playing neurodegenerative roles in astrocytes is prematurely up-regulated in an Alzheimer's model. J. Neurochem. 161: 366-382.
- Zhou, J., et al. 2023. Macrophage Gpx4 deficiency aggravates foam cell formation by regulating the expression of scavenger receptors, ABCA1, and ABCG1. Cell Biol. Int. 47: 1589-1599.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

Alexa Fluor $^{\circ}$ is a trademark of Molecular Probes, Inc., Oregon, USA