## SANTA CRUZ BIOTECHNOLOGY, INC.

# TIRAP (A-11): sc-166149



## BACKGROUND

Mammalian Toll-like receptors (TLRs) recognize conserved products of microbial metabolism and activate NF $\kappa$ B and other signaling pathways through the adapter protein MyD88. MyD88 functions as an adapter protein in the association of IL-1 receptor associated kinase (IRAK) with the IL-1 receptor. MyD88 contains a characteristic N-terminal death domain, which is essential for NF $\kappa$ B activation and an adjacent Toll/II-1R homology domain (TIR domain), which is responsible for signal transduction. TIR domain-containing adapter protein (TIRAP), also designated MAL (MyD88 adapter-like), wyatt or TLR-4 adaptor protein, is a cytoplasmic TIR-domain-containing protein that activates NF $\kappa$ B, Jun N-terminal kinase and extracellular signal-regulated kinase-1 and -2. TIRAP forms homodimers and heterodimers with MyD88. IRAK-2, but not IRAK, is required for the activation of NF $\kappa$ B by TIRAP which associates with IRAK-2 through the TIR domain. In addition, TIRAP associates with TLR-4, suggesting that it plays a role in TLR-4 signal transduction.

## **CHROMOSOMAL LOCATION**

Genetic locus: TIRAP (human) mapping to 11q24.2.

#### SOURCE

TIRAP (A-11) is a mouse monoclonal antibody raised against amino acids 1-235 representing full length TIRAP of human origin.

### PRODUCT

Each vial contains 200  $\mu g$  lgG\_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

TIRAP (A-11) is available conjugated to agarose (sc-166149 AC), 500  $\mu$ g/ 0.25 ml agarose in 1 ml, for IP; to HRP (sc-166149 HRP), 200  $\mu$ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-166149 PE), fluorescein (sc-166149 FITC), Alexa Fluor<sup>®</sup> 488 (sc-166149 AF488), Alexa Fluor<sup>®</sup> 546 (sc-166149 AF546), Alexa Fluor<sup>®</sup> 594 (sc-166149 AF594) or Alexa Fluor<sup>®</sup> 647 (sc-166149 AF647), 200  $\mu$ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor<sup>®</sup> 680 (sc-166149 AF680) or Alexa Fluor<sup>®</sup> 790 (sc-166149 AF790), 200  $\mu$ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

## **APPLICATIONS**

TIRAP (A-11) is recommended for detection of TIRAP of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for TIRAP siRNA (h): sc-42932, TIRAP shRNA Plasmid (h): sc-42932-SH and TIRAP shRNA (h) Lentiviral Particles: sc-42932-V.

Molecular Weight of TIRAP: 36 kDa.

Positive Controls: Caki-1 cell lysate: sc-2224, ES-2 cell lysate: sc-2224 or TIRAP (h): 293T Lysate: sc-114950.

#### **RECOMMENDED SUPPORT REAGENTS**

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850.

#### DATA



TIRAP (A-11): sc-1bb149. Western blot analysis of TIRAP expression in non-transfected: sc-117752 (**A**) and human TIRAP transfected: sc-114950 (**B**) 293T whole cell lysates.

#### SELECT PRODUCT CITATIONS

- Aksoy, E., et al. 2012. The p110& isoform of the kinase PI(3)K controls the subcellular compartmentalization of TLR4 signaling and protects from endotoxic shock. Nat. Immunol. 13: 1045-1054.
- Jakka, P., et al. 2018. Cytoplasmic linker protein CLIP170 negatively regulates TLR4 signaling by targeting the TLR adaptor protein TIRAP. J. Immunol. 200: 704-714.
- Rajpoot, S., et al. 2022. TIRAP-mediated activation of p38 MAPK in inflammatory signaling. Sci. Rep. 12: 5601.

#### **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

#### **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

#### **PROTOCOLS**

See our web site at www.scbt.com for detailed protocols and support products.